arXiv:math/0610505v2 [math.QA] 11 Oct 2007

Tau functions in combinatorial Bethe ansatz

Atsuo Kuniba, Reiho Sakamoto and Yasuhiko Yamada

Abstract: We introduce ultradiscrete tau functions associated with rigged configurations for $A_n^{(1)}$. They satisfy an ultradiscrete version of the Hirota bilinear equation and play a role analogous to a corner transfer matrix for the box-ball system. As an application, we establish a piecewise linear formula for the Kerov-Kirillov-Reshetikhin bijection in the combinatorial Bethe ansatz. They also lead to general N-soliton solutions of the box-ball system.

1. Introduction

The Bethe ansatz and the corner transfer matrix are methods of primary importance in analysing solvable lattice models [1]. The Bethe ansatz produces eigenvectors of row transfer matrices from solutions of the Bethe equation [2]. The corner transfer matrix method determines the one-point function from the one-dimensional sums [1]. See [3, 4, 5] and [6, 7] for some typical applications. Interestingly, both of these approaches are known to admit combinatorial versions, which have brought fruitful insights and applications into representation theory as well [8].

The combinatorial Bethe ansatz was initiated by Kerov, Kirillov and Reshetikhin (KKR) [9, 10]. They invented the object called *rigged configuration*, which serves as a combinatorial substitute for the solutions of the Bethe equation. By the KKR bijection, they are in one-to-one correspondence with the Littlewood-Richardson tableaux, or equivalently, *highest paths* which are the combinatorial analogues of the Bethe eigenvectors. As for the corner transfer matrix method, a decisive progress came with the advent of the crystal base theory [11, 12], where the one-dimensional sums are formulated as generating functions of the energy of affine crystals over paths.

Guided by a number of relevant results [13, 14, 15, 16, 17, 18], these streams have merged into the so-called $X = M$ conjecture [19, 20] for general affine Lie algebra. Here X is the one-dimensional sum in the corner transfer matrix method. For type $A_n^{(1)}$, it coincides essentially with the Kostka-Foulkes polynomial [21] for the case treated in [9, 10]. On the other hand, M is the fermionic formula (2.10) in the Bethe ansatz, which is a generating function of the charge function $c(\mu, r)$ (2.9). By now, the $X = M$ conjecture has been studied extensively and solved in several cases [22, 23, 24, 25].

During these developments, it was realized that not only the Bethe ansatz or the corner transfer matrix, but also the solvable lattice models themselves admit decent combinatorial versions. In fact, vertex models with the quantum group symmetry $U_q(A_n^{(1)})$ turned out to be the soliton cellular automata at $q = 0$ [26, 27] that had been known as the *box-ball systems* [28, 29]. Row transfer matrices in the former tend to commuting time evolutions in the latter. The finding has led to a systematic generalization of such automata [30, 31, 32], which possess fascinating features as ultradiscrete integrable systems [33]. (See the explanation under (5.10) for the ultradiscretization.) Thus it is a natural endeavor to study these automata by the combinatorial versions of the Bethe ansatz and the corner transfer matrix.

As for the Bethe ansatz, this has been done in [34, 35], which yielded the inverse scattering formalism of the box-ball systems. It turned out that rigged configurations are action-angle variables, which provide the conserved quantities or linearize the commuting time evolutions. The KKR bijection is the direct/inverse scattering (Gel'fand-Levitan) map. In particular, the mysterious combinatorial algorithm in the bijection is identified with a crystal theoretical vertex operator.

Then what about the corner transfer matrix? And this is the issue that we are going to address in this paper. From a naive point of view, one is tempted to regard the

number of balls in a quadrant of the two-dimensional time evolution pattern of the box-ball system as its candidate. We introduce such a quantity $\rho_i(p)$ (4.1) for a path p. On the other hand, the combinatorial analogue of the corner transfer matrix in the crystal base theory is the energy of affine crystals [12, 17], which is denoted by $\mathcal{E}_i(p)$ in (4.12). Our Proposition 4.6 asserts $\rho_i(p) = \mathcal{E}_i(p)$ indeed. One of the main results in this paper is Theorem 6.12, which states $\tau_i(p) = \rho_i(p) = \mathcal{E}_i(p)$. Here $\tau_i(p)$ is the piecewise linear function on the rigged configuration $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ for p:

$$\tau_i(p) = \max_{\nu \subseteq \mu} \{-c(\nu, s) - |\nu^{(i)}|\},$$
$$c(\nu, s) = \frac{1}{2} \sum_{a,b} C_{a,b} \min(\nu^{(a)}, \nu^{(b)}) - \min(\lambda, \nu^{(1)}) + \sum_a |s^{(a)}|,$$

where $(C_{ab})_{1 \le a,b \le n}$ is the Cartan matrix of A_n. $c(\nu, s)$ is the charge function appearing in the fermionic formula, and the max extends over all the subsets $(\nu^{(a)}, s^{(a)}) \subseteq (\mu^{(a)}, r^{(a)})$ of the rigged configuration. See (2.19), (2.20), (2.24) and Section 2.1 for a precise account. In short, τ_i is an ultradiscretization of a *single* summand in the fermionic formula with respect to the subsets of the rigged configuration.

An origin of this curious quantity goes back to Sato's theory of soliton equations [36]. In fact, τ_i arises as an ultradiscretization of the well known tau function for the KP hierarchy [37] under a special choice of parameters adapted to the rigged configuration. Using this fact, we show that τ_i satisfies an ultradiscrete version of the Hirota bilinear equation, which actually serves as a characterization of τ_i up to a boundary condition. We call τ_i the ultradiscrete tau function. It serves as an analogue of a corner transfer matrix in the box-ball system and bilinearize the dynamics. These features are summarized in the following table.

	Bethe ansatz	Corner transfer matrix
main combinatorial object	rigged configuration	energy in affine crystal
role in box-ball system	action-angle variable	tau function
description of dynamics	linear	bilinear

As the main consequences of Theorem 6.12, we derive a piecewise linear formula for the KKR bijection (Theorem 2.1), the solution of the initial value problem (Theorem 7.6) and the general N-soliton solution (7.21), (7.37), (7.42) for the box-ball system. Note that the quantities $\rho_i = \mathcal{E}_i$ arise from the corner transfer matrix and crystals, whereas τ_i is an explicit formula originating in the Bethe ansatz. Therefore our Theorem 6.12, i.e., $\rho_i = \mathcal{E}_i = \tau_i$ provides another connection of the two methods analogous to the $X = M$ conjecture.

The layout of the paper is as follows. In Section 2, τ_i is introduced in (2.18)–(2.20) as a piecewise linear function on rigged configurations. It is actually a member of the family $\tau_i^{(a)}$ (2.22) which obeys the recursion relation (2.23). It reflects the nested structure $sl_{n+1} \supset sl_n \supset \cdots \supset sl_2$, which will be utilized extensively. The piecewise linear formula for the KKR bijection is stated in Theorem 2.1.

In Section 3, we give the definition and the basic properties of the box-ball system.

In Section 4, we introduce ρ_i and $\mathcal{E}_i$. ρ_i in (4.1) is the number of balls in the SW quadrant in the time evolution pattern of the box-ball system. $\mathcal{E}_i$ is defined by (4.12) and (4.11), which is a sum of local energy function in the affine crystal. They are analogues of the corner transfer matrix [1] in complementary viewpoints; ρ_i originates in the box-ball system and $\mathcal{E}_i$ in the crystal base theory. They are identified in Proposition 4.6.

The piecewise linear formula for the KKR bijection (Theorem 2.1) is a consequence of the further identification $\tau_i = \rho_i = \mathcal{E}_i$ in Theorem 6.12. Sections 5 and 6 are devoted to a proof of this fact. In Section 5, τ_i is shown to emerge as an ultradiscretization of the tau functions of the KP hierarchy (Lemma 5.3) and satisfy the Hirota type bilinear equation (Proposition 5.1). The key to these results is the special choice of the parameters (5.5)–(5.9). It assures the positivity, which is vital in the ultradiscretization (Lemma 5.2). The content of this section is a refinement of the earlier analysis [26].

In Section 6, $\tau_i = \rho_i$ for $A_n^{(1)}$ is proved on the asymptotic states by induction on the rank n (Proposition 6.1 and its reduction in Proposition 6.4). From the assumption $\tau_i = \rho_i = \mathcal{E}_i$ for $A_{n-1}^{(1)}$, the scattering data is expressed in terms of tau functions (Lemma 6.6). Then we take advantage of the vertex operator formulation of the KKR bijection [34, 35] to make the induction proceed. Combined with the results in Section 5, the agreement on the asymptotic states is enough to establish the claim $\tau_i = \rho_i$ everywhere.

In Section 7, Theorem 2.1 and Theorem 6.12 are generalized to arbitrary (non-highest) states. As an application, we present the solution of the initial value problem of the box-ball system in Theorem 7.6. Our tau functions are parametrized by the conserved quantities that specify solitons. We rewrite them in several forms in (7.21), (7.37) and (7.42). They yield general N-soliton solutions of the box-ball system. Among others, our ultradiscrete tau functions are most elegantly presented in (7.42) in terms of affine crystals in the "principal picture".

Appendix A summarizes the rudiments of the crystal base theory. Appendix B illustrates the graphical rule [17] for obtaining the combinatorial R, the winding and the non-winding numbers relevant to the energy function. Appendix C recalls the combinatorial algorithm for the KKR bijection. Appendix D is the crystal theoretical reformulation of the KKR map due to [34, 35]. Appendix E is an exposition of the inverse scattering formalism of the box-ball system which supplements Section 3.

2. Ultradiscrete tau function

2.1. **Preliminary.** We summarize the basic notation used throughout the paper. For a multiset $\lambda = (\lambda_1, \dots, \lambda_k)$, we use the symbols

$$|\lambda| = \lambda_1 + \cdots + \lambda_k, \quad \ell(\lambda) = k, \tag{2.1}$$

$$\lambda_{[N]} = (\lambda_1, \dots, \lambda_N), \quad (0 \le N \le k), \tag{2.2}$$

where $\lambda_{[0]} = \emptyset$. Given two multisets $\lambda = (\lambda_1, \dots, \lambda_k)$ and $\mu = (\mu_1, \dots, \mu_m)$, we use the notation:

$$\min(\lambda, \mu) = \sum_{i=1}^{k} \sum_{j=1}^{m} \min(\lambda_i, \mu_j), \tag{2.3}$$

$$\lambda \subseteq \mu \overset{\text{def}}{\longleftrightarrow} \{\lambda_1, \dots, \lambda_k\} \subseteq \{\mu_1, \dots, \mu_m\}, \tag{2.4}$$

where $\subseteq$ accounts the multiplicity as well. For example, $\emptyset, (1,1), (1,3,1) \subseteq (1,2,1,3)$ but $(2,2) \not\subseteq (1,2,1,3)$.

2.2. **Rigged configurations.** Consider the data of the form

$$(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \dots, (\mu^{(n)}, r^{(n)})), \tag{2.5}$$

where $\mu^{(a)} = (\mu_1^{(a)}, \dots, \mu_{l_a}^{(a)}) \in (\mathbb{Z}_{\ge 1})^{l_a}$ and $r^{(a)} = (r_1^{(a)}, \dots, r_{l_a}^{(a)}) \in (\mathbb{Z}_{\ge 0})^{l_a}$ for some $l_a \ge 0$. Apart from $\mu^{(0)}$, each $(\mu^{(a)}, r^{(a)})$ is to be understood as a multiset of the pairs $(\mu_1^{(a)}, r_1^{(a)}), \dots, (\mu_{l_a}^{(a)}, r_{l_a}^{(a)})$ whose ordering does not matter. The data (2.5) is called a rigged configuration for $A_n^{(1)}$ if

$$0 \le r_i^{(a)} \le p^{(a)}_{\mu_i^{(a)}} \text{ for any pair } (\mu_i^{(a)}, r_i^{(a)}). \tag{2.6}$$

Here $p_j^{(a)}$ is called the vacancy number and defined by

$$p_j^{(a)} = E_j^{(a-1)} - 2E_j^{(a)} + E_j^{(a+1)} \quad (1 \le a \le n), \tag{2.7}$$

$$E_j^{(a)} = \sum_{k=1}^{l_a} \min(j, \mu_k^{(a)}) \ \ (0 \le a \le n), \ \ E_j^{(n+1)} = 0. \tag{2.8}$$

The array $(\mu^{(0)}, \dots, \mu^{(n)})$ is called a configuration and the nonnegative integers $r_i^{(a)}$ are called rigging. Note that $p_j^{(a)}$ and $E_j^{(a)}$ depend only on the configuration. In particular $E_\infty^{(a)} = |\mu^{(a)}|$. It is customary to arrange $\mu^{(a)}$ as $\mu^{(a)} = (\mu_1^{(a)} \ge \cdots \ge \mu_{l_a}^{(a)})$ and regard the rigged configuration as an n-tuple of Young diagrams $\mu^{(1)}, \dots, \mu^{(n)}$ where the row of length $\mu_i^{(a)}$ is assigned with the rigging $r_i^{(a)}$ subject to the condition (2.6). In this convention, we identify all the diagrams obtained by reordering the rows of equal length with different rigging. In what follows we do *not* assume $\mu_1^{(a)} \ge \cdots \ge \mu_{l_a}^{(a)}$ unless explicitly mentioned.

For a multiset with positive components λ, let $\mathrm{RC}(\lambda)$ denote the set of rigged configurations (2.5) with $\mu^{(0)} = \lambda$. Set

$$c(\mu, r) = \frac{1}{2} \sum_{a,b} C_{ab} \min(\mu^{(a)}, \mu^{(b)}) - \min(\mu^{(0)}, \mu^{(1)}) + \sum_a |r^{(a)}|, \tag{2.9}$$

where $(C_{ab})_{1 \le a,b \le n}$ is the Cartan matrix of A_n. The fermionic formula [10, 9] is obtained as the generating function:

$$M(\lambda) = \sum q^{c(\mu,r)}, \tag{2.10}$$

where the sum extends over all the rigged configurations $(\lambda, (\mu^{(1)}, r^{(1)}), \dots, (\mu^{(n)}, r^{(n)})) \in \mathrm{RC}(\lambda)$ with prescribed values for $|\mu^{(1)}|, \dots, |\mu^{(n)}|$. The sum (2.10) is arranged as $M(\lambda) = \sum_\mu q^{c(\mu,0)} \sum_r q^{\sum_{a,i} r_i^{(a)}}$, where the sum over the rigging r under the condition (2.6) yields a product of q-binomial coefficients as is well known.

2.3. **Crystals.** We recapitulate basic facts on the $A_n^{(1)}$ crystal B_l. For a general background see Appendix A. The B_l is the crystal base of the l-fold symmetric tensor representation. As the set it is given by

$$B_l = \{x = (x_1, \dots, x_{n+1}) \in (\mathbb{Z}_{\ge 0})^{n+1} \mid x_1 + \cdots + x_{n+1} = l\}. \tag{2.11}$$

The Kashiwara operators act as $\tilde{e}_i(x) = x', \tilde{f}_i(x) = x''$ with $x'_j = x_j + \delta_{i,j} - \delta_{i,j+1}$ and $x''_j = x_j - \delta_{i,j} + \delta_{i,j+1}$. Here indices are in $\mathbb{Z}_{n+1}$ and x' and x'' are to be understood as 0 unless they belong to $(\mathbb{Z}_{\ge 0})^{n+1}$. The combinatorial $R : \mathrm{Aff}(B_l) \otimes \mathrm{Aff}(B_m) \to \mathrm{Aff}(B_m) \otimes \mathrm{Aff}(B_l)$ has the form $R : x[d] \otimes y[e] \mapsto \tilde{y}[e - H(x \otimes y)] \otimes \tilde{x}[d + H(x \otimes y)]$, which are described by the piecewise linear formula [38, 26]:

$$\tilde{x}_i = x_i + Q_i(x \otimes y) - Q_{i-1}(x \otimes y), \quad \tilde{y}_i = y_i + Q_{i-1}(x \otimes y) - Q_i(x \otimes y), \tag{2.12}$$

$$Q_i(x \otimes y) = \min\{\sum_{j=1}^{k-1} x_{i+j} + \sum_{j=k+1}^{n+1} y_{i+j} \mid 1 \le k \le n+1\}, \tag{2.13}$$

$$H(x \otimes y) = \min(l, m) - Q_0(x \otimes y). \tag{2.14}$$

The energy function H here is normalized so that $0 \le H \le \min(l, m)$ and coincides with the "winding number" [17]. In general $\min(l, m) - Q_i$ is the i th winding number that counts the lines crossing x_i and x_{i+1} (Appendix B).

The element $x = (x_1, \dots, x_{n+1})$ is also denoted by a row shape semistandard tableau of length l containing the letter i x_i times and $x[d] \in \mathrm{Aff}(B_l)$ by the tableau with index

d. For example in $A_3^{(1)}$, the following stand for the same relation under R:

$$\begin{aligned}(1,2,0,1)[5]\otimes(1,0,1,0)[9]&\simeq(0,1,0,1)[8]\otimes(2,1,1,0)[6],\\ \boxed{1224}_5\otimes\boxed{13}_9&\simeq\boxed{24}_8\otimes\boxed{1123}_6.\end{aligned}\tag{2.15}$$

To save the space we use the notation:

$$a^l=\boxed{a\cdots a}\in B_l,\quad u_l=1^l=\boxed{1\cdots 1}\in B_l.\tag{2.16}$$

Setting

$$B_l^{\geq a+1}=\{(x_1,\dots,x_{n+1})\in B_l\mid x_1=\cdots=x_a=0\}\quad(0\leq a\leq n),$$

we have

$$B_l=B_l^{\geq 1}\supset B_l^{\geq 2}\supset\cdots\supset B_l^{\geq n+1}=\{(n+1)^l\}\tag{2.17}$$

as sets. We will need to consider the crystals not only for $A_n^{(1)}$ but also for the nested family $A_0^{(1)},A_1^{(1)},\dots,A_{n-1}^{(1)}$. In such a circumstance we realize the crystal B_l for $A_{n-a}^{(1)}$ $(0\leq a\leq n)$ on the set $B_l^{\geq a+1}$ with the Kashiwara operators $\tilde{e}_i,\tilde{f}_i$ $(a\leq i\leq n)$. In this convention the highest element with respect to A_{n-a} is $(a+1)^l\in B_l^{\geq a+1}$.

Let

$$\mathcal{P}_+(\lambda)=\{p\in B_{\lambda_1}\otimes\cdots\otimes B_{\lambda_L}\mid\tilde{e}_ip=0,\ 1\leq i\leq n\}.$$

be the set of highest elements (paths) with respect to A_n. The bijection [9, 10] between $\mathrm{RC}(\lambda)$ and the Littlewood-Richardson tableaux is translated to the one between $\mathrm{RC}(\lambda)$ and $\mathcal{P}_+(\lambda)$. We call the resulting map the KKR bijection. See Appendix C for an exposition of the algorithm and Appendix D for the recent reformulation as the crystal theoretical vertex operator [34, 35]. In particular, there is a nested structure with respect to the rank in the sense that if $(\mu^{(0)},(\mu^{(1)},r^{(1)}),\dots,(\mu^{(n)},r^{(n)}))$ is a rigged configuration for $A_n^{(1)}$, so is $(\mu^{(a)},(\mu^{(a+1)},r^{(a+1)}),\dots,(\mu^{(n)},r^{(n)}))$ for $A_{n-a}^{(1)}$. Moreover, the KKR bijection sends the latter to a highest path in $B^{\geq a+1}_{\mu_1^{(a)}}\otimes\cdots\otimes B^{\geq a+1}_{\mu_{l_a}^{(a)}}$.

2.4. **Piecewise linear formula for KKR bijection.** We use the notation defined in Section 2.1. Given a rigged configuration $(\mu^{(0)},(\mu^{(1)},r^{(1)}),\dots,(\mu^{(n)},r^{(n)}))$, we introduce the ultradiscrete tau functions $\tau_0(\lambda),\tau_1(\lambda),\dots,\tau_{n+1}(\lambda)$ for $\lambda\subseteq\mu^{(0)}$ as follows:

$$\tau_0(\lambda)=\tau_{n+1}(\lambda)-|\lambda|,\tag{2.18}$$

$$\tau_d(\lambda)=\max_{\nu\subseteq\mu}\{-c(\nu,s)-|\nu^{(d)}|\}\quad(1\leq d\leq n+1,\ |\nu^{(n+1)}|=0),\tag{2.19}$$

$$\begin{aligned}-c(\nu,s)=&\min(\lambda,\nu^{(1)})+\min(\nu^{(1)},\nu^{(2)})+\cdots+\min(\nu^{(n-1)},\nu^{(n)})\\&-\min(\nu^{(1)},\nu^{(1)})-\min(\nu^{(2)},\nu^{(2)})-\cdots-\min(\nu^{(n)},\nu^{(n)})\\&-|s^{(1)}|-\cdots-|s^{(n)}|.\end{aligned}\tag{2.20}$$

In (2.19), max is taken over $\nu=(\nu^{(1)},\dots,\nu^{(n)})$, where the components are independently chosen under the condition $\nu^{(1)}\subseteq\mu^{(1)},\dots,\nu^{(n)}\subseteq\mu^{(n)}$. The array $s=(s^{(1)},\dots,s^{(n)})$ denotes the set of the riggings $s^{(1)}\subseteq r^{(1)},\dots,s^{(n)}\subseteq r^{(n)}$ that are paired with the chosen $\nu^{(1)},\dots,\nu^{(n)}$ as $\{(\nu_i^{(a)},s_i^{(a)})\}\subseteq\{(\mu_i^{(a)},r_i^{(a)})\}$. The quantity $c(\nu,s)$ in (2.20) is obtained from $c(\mu,r)$ (2.9) by replacing $(\mu,r)=(\mu^{(0)},(\mu^{(1)},r^{(1)}),\dots,(\mu^{(n)},r^{(n)}))$ with $(\nu,s)=(\lambda,(\nu^{(1)},s^{(1)}),\dots,(\nu^{(n)},s^{(n)}))$. Apart from $\nu^{(a)}\subseteq\mu^{(a)}$, there is *no* further constraint on $|\nu^{(1)}|,\dots,|\nu^{(n)}|$ and it is *not* required that the data $(\lambda,(\nu^{(1)},s^{(1)}),\dots,(\nu^{(n)},s^{(n)}))$ to be a rigged configuration for $A_n^{(1)}$. Since the max (2.19) includes the trivial case $\forall\nu^{(a)}=\emptyset$, the quantities $\tau_1(\lambda),\dots,\tau_{n+1}(\lambda)$ are nonnegative integers. Note that $\tau_{n+1}(\mu^{(0)})=\max\{-c(\nu,s)\}$ in (2.19) may be viewed as an ultradiscretization of the *single* summand $q^{c(\mu,r)}$ in the fermionic formula (2.10) with respect to the subsets $(\nu,s)\subseteq(\mu,r)$. See also (5.11).

Theorem 2.1. *Let the image of the rigged configuration* $(\mu^{(0)},(\mu^{(1)},r^{(1)}),\dots,(\mu^{(n)},r^{(n)}))$ *under the KKR bijection be the highest path* $p_1\otimes\cdots\otimes p_L\in\mathcal{P}_+(\mu^{(0)})$. *Then* $p_k=(x_1,\dots,x_{n+1})\in B_{\mu_k^{(0)}}$ *is expressed as*

$$x_d=\tau_{k,d}-\tau_{k-1,d}-\tau_{k,d-1}+\tau_{k-1,d-1}, \tag{2.21}$$

where $\tau_{k,d}=\tau_d((\mu_1^{(0)},\dots,\mu_k^{(0)}))$.

Note that (2.18) ensures $x_1+\cdots+x_{n+1}=\mu_k^{(0)}$.

Due to the nested structure of the KKR bijection with respect to the rank [34], Theorem 2.1 is also stated as a family of relations corresponding to $sl_{n+1}\supset sl_n\supset\cdots\supset sl_2$. To do so, we introduce the family of ultradiscrete tau functions $\{\tau_d^{(a)}(\lambda)\mid 0\le a\le n-1,\ a\le d\le n+1,\ \lambda\subseteq\mu^{(a)}\}$ by $\tau_a^{(a)}(\lambda)=\tau_{n+1}^{(a)}(\lambda)-|\lambda|$ and

$$\begin{aligned}\tau_d^{(a)}(\lambda)=\max\{&\min(\lambda,\nu^{(a+1)})+\min(\nu^{(a+1)},\nu^{(a+2)})+\cdots+\min(\nu^{(n-1)},\nu^{(n)})\\&-\min(\nu^{(a+1)},\nu^{(a+1)})-\min(\nu^{(a+2)},\nu^{(a+2)})-\cdots-\min(\nu^{(n)},\nu^{(n)})\\&-|s^{(a+1)}|-|s^{(a+2)}|-\cdots-|s^{(n)}|-|\nu^{(d)}|\}\quad(a+1\le d\le n+1),\end{aligned} \tag{2.22}$$

where $|\nu^{(n+1)}|=0$ as before. The max is taken over the independent choices $\nu^{(a+1)}\subseteq\mu^{(a+1)},\dots,\nu^{(n)}\subseteq\mu^{(n)}$. The subsets of the riggings $s^{(a+1)}\subseteq r^{(a+1)}$, …, $s^{(n)}\subseteq r^{(n)}$ are those paired with the chosen $\nu^{(a+1)},\dots,\nu^{(n)}$ as before. The previously introduced tau function $\tau_d(\lambda)$ (2.19) is equal to $\tau_d^{(0)}(\lambda)$. Now Theorem 2.1 is rephrased as

Theorem 2.2. *Given a rigged configuration* $(\mu^{(0)},(\mu^{(1)},r^{(1)}),\dots,(\mu^{(n)},r^{(n)}))$ *and* $0\le a\le n-1$, *let the image of* $(\mu^{(a)},(\mu^{(a+1)},r^{(a+1)}),\dots,(\mu^{(n)},r^{(n)}))$ *under the KKR bijection be the* A_{n-a} *highest path* $p_1\otimes\cdots\otimes p_{l_a}\in B^{\ge a+1}_{\mu_1^{(a)}}\otimes\cdots\otimes B^{\ge a+1}_{\mu_{l_a}^{(a)}}$. *Then* $p_k=(x_{a+1},x_{a+2},\dots,x_{n+1})$ *is expressed as*

$$x_d=\tau_{k,d}^{(a)}-\tau_{k-1,d}^{(a)}-\tau_{k,d-1}^{(a)}+\tau_{k-1,d-1}^{(a)},$$

where $\tau_{k,d}^{(a)}=\tau_d^{(a)}((\mu_1^{(a)},\dots,\mu_k^{(a)}))$.

Again, $x_{a+1}+\cdots+x_{n+1}=\mu_k^{(a)}$ is evident by the construction. For a proof of Theorem 2.1, see Section 4.4.

The tau functions (2.22) are the solution of the recursion relation with respect to the rank:

$$\tau_d^{(a)}(\lambda)=\max_{\nu\subseteq\mu^{(a+1)}}\{\min(\lambda,\nu)-\min(\nu,\nu)-|s|+\tau_d^{(a+1)}(\nu)\} \tag{2.23}$$

for $0\le a\le n-1$, $a+1\le d\le n+1$ with the convention $\tau_a^{(a)}(\lambda)=\tau_{n+1}^{(a)}(\lambda)-|\lambda|$ and the initial condition $\tau_{n+1}^{(n)}(\lambda)=0$, $\tau_n^{(n)}(\lambda)=-|\lambda|$. The rigging s is the subset of $r^{(a+1)}$ paired with the chosen ν.

Lemma 2.3. $\tau_d^{(a)}(\emptyset)=0$ *for any* $0\le a\le n-1$ *and* $a+1\le d\le n+1$.

Proof. It suffices to prove $a=0$ case. When $\lambda=\emptyset$, (2.19) becomes $\tau_d(\emptyset)=-\min_{\nu\subseteq\mu}\{\tilde{c}(\nu)+|s^{(1)}|+\cdots+|s^{(n)}|+|s^{(d)}|\}$, where $\tilde{c}(\nu)$ is given by (see (2.9))

$$\tilde{c}(\nu)=\frac{1}{2}\sum_{a,b}C_{a,b}\min(\nu^{(a)},\nu^{(b)})=\frac{1}{2}\sum_{a,b}C_{a,b}\sum_{i,j}\min(i,j)m_i^{(a)}m_j^{(b)},$$

where $m_i^{(a)}$ is the number of k such that $\nu_k^{(a)}=i$. This is a positive definite quadratic form whose minimum is 0 at $\forall m_j^{(a)}=0$. The other part $|s^{(1)}|+\cdots+|s^{(n)}|+|s^{(d)}|$ appearing in $\tau_d(\emptyset)$ also attains the minimum 0 simultaneously at this point. □

Let the image of the rigged configuration $(\mu^{(0)},(\mu^{(1)},r^{(1)}),\dots,(\mu^{(n)},r^{(n)}))$ under the KKR bijection be the highest path $p_1\otimes\cdots\otimes p_L\in\mathcal{P}_+(\mu^{(0)})\subset B_{\mu_1^{(0)}}\otimes\cdots\otimes B_{\mu_L^{(0)}}$. In what follows we will also write

$$\tau_i(\lambda)=\tau_{k,i}=\tau_i(p_1\otimes\cdots\otimes p_k)\ \text{ for }\ \lambda=\mu^{(0)}_{[k]}=(\mu^{(0)}_1,\dots,\mu^{(0)}_k)\quad(1\le k\le L). \tag{2.24}$$

Concerning the notation $\tau_i(p_1\otimes\cdots\otimes p_k)$, a remark is in order. Any highest path $p_1\otimes\cdots\otimes p_k$ can be extended to a longer one $p_1\otimes\cdots\otimes p_k\otimes p_{k+1}\otimes\cdots\otimes p_L$ in which $p_{k+1}\otimes\cdots\otimes p_L$ is not unique. Suppose that $(\mu^{(0)},(\mu^{(1)},r^{(1)}),\dots,(\mu^{(n)},r^{(n)}))$ and $(\mu'^{(0)},(\mu'^{(1)},r'^{(1)}),\dots,(\mu'^{(n)},r'^{(n)}))$ are two rigged configurations corresponding to such extensions of $p_1\otimes\cdots\otimes p_k$, and let $\tau_i(p_1\otimes\cdots\otimes p_k)$ and $\tau'_i(p_1\otimes\cdots\otimes p_k)$ be the associated tau functions in the sense of (2.24). Then $\tau_i(p_1\otimes\cdots\otimes p_k)=\tau'_i(p_1\otimes\cdots\otimes p_k)$ will be guaranteed by Theorem 4.9. Note however that they are different as the piecewise linear expressions as in (2.18)–(2.20). By the reason, we will always mention the rigged configurations relevant to $p_1\otimes\cdots\otimes p_k$.

Example 2.4. Consider the highest path $p=11112221322433\in B_1^{\otimes L}$ of length $L=14$, where we have omitted the symbol $\otimes$. The corresponding rigged configuration is depicted in Example C.2. Thus we set

$$\mu^{(0)}=(1^{14}),\ \ \mu^{(1)}=(4,3,2),\ \ \mu^{(2)}=(3,1),\ \ \mu^{(3)}=(1),$$
$$r^{(1)}=(0,2,3),\ \ \ r^{(2)}=(1,0),\ \ \ r^{(3)}=(0).$$

The associated tau function $\tau_{k,i}$ takes the following values.

k	1	2	3	4	5	6	7	8	9	10	11	12	13	14
$\tau_{k,1}$	0	0	0	0	0	0	0	1	2	3	4	6	8	10
$\tau_{k,2}$	0	0	0	0	1	2	3	4	5	7	9	11	13	15
$\tau_{k,3}$	0	0	0	0	1	2	3	4	6	8	10	12	15	18
$\tau_{k,4}$	0	0	0	0	1	2	3	4	6	8	10	13	16	19

The choices of the subsets $\nu=(\nu^{(1)},\nu^{(2)},\nu^{(3)})$ that attain these values for $\tau_{k,4}=\max_{\nu\subseteq\mu}\{\cdots\}$ in (2.19) are as follows.

k	ν
$1,2,3$	A
4	A,B
$5,6,7$	B
8	B,C
$9,10$	C
11	C,D
$12,13,14$	D

Here $A,B,C,D\subseteq\mu=(\mu^{(1)},\mu^{(2)},\mu^{(3)})$ are given by

$$A=(\emptyset,\emptyset,\emptyset),$$
$$B=((4),\emptyset,\emptyset),\ ((4),(1),\emptyset),\ ((4),(1),(1)),$$
$$C=((4,2),(3),\emptyset),\ ((4,2),(1),\emptyset),\ ((4,2),(3),(1)),\ ((4,2),(1),(1)),\ ((4,2),(3,1),(1)),$$
$$D=((4,3,2),(3,1),(1))=(\mu^{(1)},\mu^{(2)},\mu^{(3)}).$$

The case $k=0$ enforces the choice $\forall\nu^{(a)}=\emptyset$ in agreement with Lemma 2.3. In the other extreme case $k=L$, the full choice $\nu=\mu$ is the consequence of the general result in Remark 6.14. In general the maximum attaining ν for $\tau_i(\lambda)=\max_{\nu\subseteq\mu}\{\cdots\}$ gradually grows with λ. The above p will be investigated further in Examples E.1 and E.4.

3. Box-ball system

3.1. **Conventional formulation.** Consider the tensor product $B_{\lambda_1} \otimes B_{\lambda_2} \otimes \cdots \otimes B_{\lambda_L}$. Its elements are called states. We regard each component $(x_1, \ldots, x_{n+1}) \in B_l$ as a capacity l box containing x_i balls with color i for $2 \le i \le n+1$. On the other hand the letter 1 is to be interpreted as a vacancy. Thus x_1 represents the empty space in the box. A state represents an array of boxes with capacity $\lambda_1, \ldots, \lambda_L$ containing balls of colors $2, 3, \ldots, n+1$.

We define the time evolution $T_l(p) = p'_1 \otimes \cdots \otimes p'_L$ of a state $p = p_1 \otimes \cdots \otimes p_L$ by

$$u_l[0] \otimes p_1[0] \otimes \cdots \otimes p_L[0] \simeq p'_1[-d_1] \otimes \cdots \otimes p'_L[-d_L] \otimes v_l[d_1 + \cdots + d_L] \tag{3.1}$$

under the isomorphism $\mathrm{Aff}(B_l) \otimes (\mathrm{Aff}(B_{\lambda_1}) \otimes \cdots \otimes \mathrm{Aff}(B_{\lambda_L})) \simeq (\mathrm{Aff}(B_{\lambda_1}) \otimes \cdots \otimes \mathrm{Aff}(B_{\lambda_L})) \otimes \mathrm{Aff}(B_l)$. Here $v_l \in B_l$ and d_i are uniquely determined by (2.12)–(2.14). We set

$$E_l(p) = e_1 + \cdots + e_L, \quad e_j = \min(\lambda_j, l) - d_j, \tag{3.2}$$

which has the property $E_l(p \otimes u_k) = E_l(p)$ for any k and l.

It is known [31, 27, 26] that T_l is weight preserving, the commutativity $T_l T_k = T_k T_l$ is valid and $E_l(p)$ is a conserved quantity, i.e., $E_l(T_k(p)) = E_l(p)$ for any k and l, provided that $p_j = u_{\lambda_j}$ for $L' \le j \le L$ with sufficiently large $L - L'$. The proof of these facts is based on the Yang-Baxter equation of the combinatorial R (Proposition A.1) and the property:

$$v_l = u_l \ \text{ if } p_j = u_{\lambda_j} \text{ for } L' \le j \le L \text{ with sufficiently large } L - L'. \tag{3.3}$$

T_l stabilizes for $l \gg 1$, which will be denoted by T_∞.

Since each d_j is the winding number (2.14), $E_l(p)$ is the sum of the non-winding number e_j. In particular for $l = \infty$, e_j is equal to the number of balls $x_2 + \cdots + x_{n+1}$ in the j th box $p_j = (x_1, x_2, \ldots, x_{n+1}) \in B_{\lambda_j}$. Therefore we find

$$E_\infty(p) = \text{number of balls contained in } p. \tag{3.4}$$

In the terminology of solvable lattice models, E_l is the energy associated with a row transfer matrix. It should not be confused with another energy $\mathcal{E}_i$ (4.12) relevant to the corner transfer matrix. Their relation is given in Proposition 4.8. The conserved quantity E_l will be evaluated explicitly for highest states in Proposition 6.15 and for general states in Proposition 7.7.

Example 3.1. The time evolution of the top row p under T_∞, i.e., $T_\infty(p)$, $T_\infty^2(p)$, $T_\infty^3(p)$ are listed downward. The frame of the semistandard tableaux and the symbol $\otimes$ are omitted.

11	122	2	1333	1	1	4	1	1	1	1	1	1	1	1	1	1
11	111	1	1222	3	3	3	4	1	1	1	1	1	1	1	1	1
11	111	1	1111	2	2	2	3	4	3	3	1	1	1	1	1	1
11	111	1	1111	1	1	1	2	3	2	2	4	3	3	1	1	1

The conserved quantities are given by $E_1(p) = 3, E_2(p) = 5$ and $E_l(p) = 7$ for $l \ge 3$.

The time evolution T_∞ can be calculated by a simple prescription [26]. We introduce a map $L_i \, (2 \le i \le n+1)$ by

$$\begin{aligned} L_i : \mathbb{Z}_{\ge 0} \times B_l &\longrightarrow B_l \times \mathbb{Z}_{\ge 0} \\ (m, y) &\longmapsto (y', m'), \end{aligned} \tag{3.5}$$

where m' and $y' = (y'_1, \ldots, y'_{n+1})$ are determined from m and $y = (y_1, \ldots, y_{n+1})$ by

$$m' = y_i + (m - y_1)_+, \quad y'_j = \begin{cases} y_i + (y_1 - m)_+ & \text{if } j = 1, \\ \min(m, y_1) & \text{if } j = i, \\ y_j & \text{otherwise}, \end{cases} \tag{3.6}$$

where $(m)_+ = \max(m, 0)$. L_i may be viewed as the interaction of the box B_l with the carrier that contains m balls of color i. The carrier drops as many balls as possible into the empty space y_1 and picks away all the color i balls that were originally in the box. Using L_i, we introduce the operators K_i $(2 \le i \le n+1)$ that sends a state to another as follows.

$$K_i(p_1 \otimes p_2 \otimes \cdots) = p_1' \otimes p_2' \otimes \cdots,$$
$$L_i((m_j, p_j)) = (p_j', m_{j+1}) \text{ for } j \ge 0, \ (m_0 = 0).$$

The latter relation is applied successively for $j = 0, 1, 2, \ldots$, determining all the p_j''s. In other words the operator K_i attaches an empty carrier to the left of the state and sends it to the right, by which the color i balls are moved to the right according to the local interaction rule L_i.

Proposition 3.2 ([26])**.** *The time evolution T_∞ admits the factorization:*

$$T_\infty = K_2 K_3 \cdots K_{n+1}.$$

Example 3.3. For p in Example 3.1, $K_4(p), K_3K_4(p)$ and $K_2K_3K_4(p) = T_\infty(p)$ are given.

11	122	2	1333	1	1	4	1	1	1	1	1	1	1	1	1	1
11	122	2	1333	1	1	1	4	1	1	1	1	1	1	1	1	1
11	122	2	1111	3	3	3	4	1	1	1	1	1	1	1	1	1
11	111	1	1222	3	3	3	4	1	1	1	1	1	1	1	1	1

Remark 3.4. Suppose $p_j = u_{\lambda_j}$ for $1 \le j \le k$ in a state $p = p_1 \otimes \cdots \otimes p_L$. Then Proposition 3.2 tells that in the state $T_\infty(p) = p_1' \otimes \cdots \otimes p_L'$, $p_j' = u_{\lambda_j}$ is valid for $1 \le j \le k+1$.

3.2. **Bethe ansatz.** Highest states in $B_{\lambda_1} \otimes \cdots \otimes B_{\lambda_L}$ are in one to one correspondence with rigged configurations $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ with $\mu^{(0)} = (\lambda_1, \ldots, \lambda_L)$ by the KKR bijection. Suppose L is sufficiently large. If a state $p = p_1 \otimes \cdots \otimes p_L$ is highest and $p_k = u_{\lambda_k}$ for $k \gg 1$, so is its time evolution $T_l(p)$. Thus the box-ball system induces the time evolution on the associated rigged configurations. For such states, $E_j^{(0)}$ (2.8) and the vacancy number $p_j^{(1)}$ are sufficiently large, and one can increase the color 1 rigging $r_i^{(1)}$ without violating the condition (2.6).

Proposition 3.5 ([34], Proposition 2.6)**.** *Let $p = p_1 \otimes \cdots \otimes p_L \in \mathcal{P}_+(\mu^{(0)})$ be the image of the rigged configuration $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ under the KKR bijection. Assume that $v_l = u_l$ in (3.1) and set $r_i^{'(1)} = r_i^{(1)} + \min(l, \mu_i^{(1)})$.*
Then $(\mu^{(0)}, (\mu^{(1)}, r^{'(1)}), (\mu^{(2)}, r^{(2)}), \ldots, (\mu^{(n)}, r^{(n)}))$ is a rigged configuration and corresponds to the highest state $T_l(p) \in \mathcal{P}_+(\mu^{(0)})$.

This is proved from the definition of the time evolution (3.1) and Lemma C.3. The time evolution T_l in this paper corresponds to the $a = 1$ case of $T_l^{(a)}$ considered in [34]. In this sense the rigged configurations are the action-angle variables of the box-ball system which linearize the original nonlinear dynamics (3.1). Moreover it is clear that all the $T_l(p)$ are the same if $l \ge \max \mu^{(1)}$.

Example 3.6. The rigged configuration corresponding to $T_\infty^t(p)$ $(t = 0, 1, 2, 3)$ in Example 3.1 (apart from $\mu^{(0)}$).



The length of each row is $\mu_i^{(a)}$ and the numbers on its right and left are the rigging $r_i^{(a)}$ and the vacancy number $p^{(a)}_{\mu_i^{(a)}}$, respectively. (Vacancy numbers are exhibited here for a check of (2.6).)

The Bethe ansatz produces transfer matrix eigenvectors from solutions to Bethe equations. The KKR bijection is its combinatorial version in the sense that the former is replaced by highest states and the latter by rigged configurations. Thus we see that the combinatorial Bethe ansatz provides a linearization scheme, or equivalently, an inverse scattering method of the box-ball system [34]. See Appendix E for a further exposition combined with the vertex operator formalism of the KKR bijection.

4. Corner transfer matrix

4.1. **Number of balls in the SW quadrant.** Let $p = p_1 \otimes \cdots \otimes p_L$ be a state and write its time evolution as $T^t_\infty(p_1 \otimes \cdots \otimes p_L) = p_1^t \otimes \cdots \otimes p_L^t$, with $p_j^t = (x^t_{j,1}, x^t_{j,2}, \ldots, x^t_{j,n+1}) \in B_{\lambda_j}$. We do not assume that p is highest. For $0 \le k \le L$ and $1 \le d \le n+1$, we define the function $\rho_{k,d}(p) \in \mathbb{Z}_{\ge 0}$ by $(\rho_{0,d}(p) = 0)$

$$\rho_{k,d}(p) = \sum_{j=1}^{k} (x^0_{j,2} + \cdots + x^0_{j,d}) + \sum_{t \ge 1} \sum_{j=1}^{k} (x^t_{j,2} + \cdots + x^t_{j,n+1}). \tag{4.1}$$

Here the second term is finite due to Remark 3.4. In fact the double sum may well be replaced by $\sum_{t=1}^{k-1} \sum_{j=t+1}^{k}$ only where the nonzero contributions are contained. This region is depicted as the SW quadrant of the time evolution pattern like Example 3.1.



The first term in (4.1) is the number of balls of color $2, 3, \ldots, d$ contained in the top row, which is the truncation $p_1 \otimes \cdots \otimes p_k$ of the state p. The second term counts the balls of all colors $2, \ldots, n+1$ within the hatched domain. By the definition, $\rho_{k,n+1}$ is the total number of balls within $p_1 \otimes \cdots \otimes p_k$ and the SW quadrant beneath it. Thus

$$\rho_{k,1}(p) = \rho_{k,n+1}(T_\infty(p)) \tag{4.2}$$

holds. Note that $\rho_{k,d}(p)$ is independent of $p_{k+1}, p_{k+2}, \ldots, p_L$. In this regard, we will also use the notation

$$\rho_d(p_1 \otimes \cdots \otimes p_k) = \rho_{k,d}(p). \tag{4.3}$$

From Remark 3.4 it follows that

$$\rho_d(u_l \otimes p_1 \otimes \cdots \otimes p_k) = \rho_d(p_1 \otimes \cdots \otimes p_k) \tag{4.4}$$

for any l.

The above picture reminds us of Baxter's corner transfer matrix (CTM) in solvable lattice models [1]. In fact $\rho_{k,d}$ serves its ultradiscrete analogue adapted to the box-ball system as we will see below.

Example 4.1. For p in Example 3.1, $\rho_{k,d}(p)$ takes the following values.

k	1	2	3	4	5	6	7	8
$\rho_{k,1}$	0	0	0	3	5	7	9	12
$\rho_{k,2}$	0	2	3	6	8	10	12	15
$\rho_{k,3}$	0	2	3	9	11	13	15	18
$\rho_{k,4}$	0	2	3	9	11	13	16	19

4.2. **Bilinearization of box-ball system.** By the definition, the k th component $p_k = (x_1, \ldots, x_{n+1}) \in B_{\lambda_k}$ in a state $p = p_1 \otimes \cdots \otimes p_L$ is expressed as

$$x_d = \rho_{k,d} - \rho_{k-1,d} - \rho_{k,d-1} + \rho_{k-1,d-1} \quad (1 \le d \le n+1), \tag{4.5}$$

where $\rho_{k,d} = \rho_{k,d}(p)$ for $1 \le d \le n+1$ and the extra one $\rho_{k,0}(p)$ is specified by

$$\rho_{k,0}(p) = \rho_{k,n+1}(p) - (\lambda_1 + \cdots + \lambda_k) \tag{4.6}$$

so as to satisfy $x_1 + \cdots + x_{n+1} = \lambda_k$. The formula (4.5) may be viewed, in a certain sense, as an ultradiscrete analogue of the Baxter formula (eq.(13.1.12) in [1]): $\langle \sigma_1 \rangle = \mathrm{Tr}(SABCD)/\mathrm{Tr}(ABCD)$ for one point function in terms of CTMs.

We use the notation

$$\overline{\rho}_{k,d} = \rho_{k,d}(T_\infty(p)). \tag{4.7}$$

Thus (4.2) reads

$$\rho_{k,1} = \overline{\rho}_{k,n+1}. \tag{4.8}$$

Proposition 4.2. *For $2 \le d \le n+1$ the following relation holds:*

$$\overline{\rho}_{k,d-1} + \rho_{k-1,d} = \max(\overline{\rho}_{k,d} + \rho_{k-1,d-1},\ \overline{\rho}_{k-1,d-1} + \rho_{k,d} - \lambda_k). \tag{4.9}$$

A similar fact has been shown in [26].

Proof. In the time evolution $T_\infty = K_2 K_3 \cdots K_{n+1}$ (Proposition 3.2), let us calculate the effect of the operator K_d on the k th box $p_k = (x_1, \ldots, x_{n+1}) \in B_{\lambda_k}$ in $K_{d+1} \cdots K_{n+1}(p)$. In the following, the fact that color d balls are touched only by K_d is taken into account. Suppose that the carrier contains m and m' balls with color d just before and after the interaction L_d (3.5). In (3.6) we are to set

$$\begin{aligned} m' &= \sum_{j=1}^{k} (\rho_{j,d} - \rho_{j-1,d} - \rho_{j,d-1} + \rho_{j-1,d-1}) - (\rho \to \overline{\rho}) \\ &= (\rho_{k,d} - \rho_{k,d-1}) - (\overline{\rho}_{k,d} - \overline{\rho}_{k,d-1}), \\ m &= m'|_{k \to k-1}, \\ y_d &= x_d = \rho_{k,d} - \rho_{k-1,d} - \rho_{k,d-1} + \rho_{k-1,d-1}, \end{aligned}$$

where we have used (4.5). As for the empty space y_1 concerning L_d in (3.5), we show that it is given by

$$y_1 = \lambda_k + \overline{\rho}_{k,d} - \overline{\rho}_{k-1,d} - \rho_{k,d} + \rho_{k-1,d} \quad (2 \le d \le n+1) \tag{4.10}$$

by induction on d in the decreasing order $d = n+1, n, \ldots, 2$. In so doing, the bilinear relation (4.9) will be established simultaneously.

For $d = n+1$, (4.10) coincides with x_1 in (4.5) by (4.6) and (4.8), hence it is correct. Then the relation $m' = y_d + (m - y_1)_+$ (3.6) leads to (4.9). The new empty space is determined from $y_1' = y_d + (y_1 - m)_+ = m' + y_1 - m$ and is equal to

$$\lambda_k + \overline{\rho}_{k,d-1} - \overline{\rho}_{k-1,d-1} - \rho_{k,d-1} + \rho_{k-1,d-1}.$$

This coincides with (4.10) with d replaced by $d-1$, making the induction proceed. □

The relation (4.9) is an ultradiscrete analogue of the Hirota bilinear equation. In view of (4.8), it determines $\rho_{k-1,1}, \rho_{k-1,2}, \ldots, \rho_{k-1,n+1}$ successively from $\{\overline{\rho}_{k-1,d}, \overline{\rho}_{k,d}, \rho_{k,d} \mid 1 \le d \le n+1\}$. Thus all the $\rho_{k,d}(T^t_\infty(p))$ are fixed uniquely from the data at sufficiently large t and k. Then the local states are specified by (4.5). In this sense the ultradiscrete CTM ρ_d achieves a bilinearization of the dynamics of the box-ball system.

4.3. **Relation to energy function.** Let $p \in B_{\lambda_1} \otimes \cdots \otimes B_{\lambda_L}$ be any element which is not necessarily highest. For $1 \le k \le L$, we introduce the sum:

$$\mathcal{E}^\vee_i(p_1 \otimes \cdots \otimes p_k) = \sum_{1 \le j < m \le k} Q_i(p_j \otimes p_m^{(j+1)}) \quad (1 \le i \le n+1), \tag{4.11}$$

where Q_i is the i th non-winding number (2.13) with the convention $Q_{n+1} = Q_0$. The element $p_m^{(j+1)}$ is defined by sending p_m to the left by applying the combinatorial R successively as

$$\begin{aligned} p_j \otimes p_{j+1} \otimes \cdots \otimes p_{m-1} \otimes p_m &\simeq p_j \otimes p_{j+1} \otimes \cdots \otimes p_m^{(m-1)} \otimes p'_{m-1} \simeq \cdots \\ &\simeq p_j \otimes p_{j+1} \otimes p_m^{(j+2)} \otimes \cdots \otimes p'_{m-1} \\ &\simeq p_j \otimes p_m^{(j+1)} \otimes p'_{j+1} \otimes \cdots \otimes p'_{m-1}. \end{aligned}$$

We understand that (4.11) is 0 for $k = 0, 1$. Using $\mathcal{E}^\vee_i$ we define the i th energy $\mathcal{E}_i$ by

$$\mathcal{E}_i(p_1 \otimes \cdots \otimes p_k) = \mathcal{E}^\vee_i(u_\infty \otimes p_1 \otimes \cdots \otimes p_k) \quad (1 \le i \le n+1), \tag{4.12}$$

where u_∞ actually means u_l with sufficiently large l. $\mathcal{E}_i$ does not depend on such l. In fact, from the graphical rule in Appendix B, we find $Q_i(u_l \otimes x) = x_2 + x_3 + \cdots + x_i$ if $x = (x_1, \ldots, x_{n+1})$ and $x_1 + \cdots + x_{n+1} \le l$ is satisfied. Thus writing $p_j^{(1)} = (x_{j,1}, \ldots, x_{j,n+1})$, (4.12) is split into the boundary and the bulk parts as

$$\mathcal{E}_i(p_1 \otimes \cdots \otimes p_k) = \sum_{j=1}^{k} (x_{j,2} + \cdots + x_{j,i}) + \mathcal{E}^\vee_i(p_1 \otimes \cdots \otimes p_k). \tag{4.13}$$

In particular, one has $\mathcal{E}_i(p_1 \otimes \cdots \otimes p_k) = \mathcal{E}^\vee_i(p_1 \otimes \cdots \otimes p_k)$ if $p_1 \otimes \cdots \otimes p_k$ is highest. We warn that the quantity usually called energy [17, 20] is $-\mathcal{E}_{n+1}$ or $-\mathcal{E}^\vee_{n+1}$ up to an additive constant. In what follows, whenever the notation u_∞ is used, it should be understood as u_l with sufficiently large l and the relevant quantity is independent of such l.

To the relation $x \otimes y \simeq \tilde{y} \otimes \tilde{x}$ with $e = Q_i(x \otimes y)$, we assign the diagram

(4.14)



where the suppressed i is to be mentioned nearby if necessary.

Let $\sigma((x_1, x_2, \ldots, x_{n+1})) = (x_2, x_3, \ldots, x_1)$ be the Dynkin diagram automorphism acting on B_l decreasing the tableau letters cyclically by one. We extend it naturally to the tensor product by $\sigma(p_1 \otimes \cdots \otimes p_k) = \sigma(p_1) \otimes \cdots \otimes \sigma(p_k)$. Since the combinatorial R commutes with σ, the i th non-winding number has the properties similar to the $i = 0$ case. In particular, under the Yang-Baxter relation



the equalities $a + b = e + f$ and $b + c = d + e$ hold. In fact, suppose the figure corresponds to $\mathrm{Aff}(B_k) \otimes \mathrm{Aff}(B_l) \otimes \mathrm{Aff}(B_m) \to \mathrm{Aff}(B_m) \otimes \mathrm{Aff}(B_l) \otimes \mathrm{Aff}(B_k)$ for some k, l and m.

If $i = 0$ for instance, the associated non-winding number Q_0 is related to H via (2.14), therefore by setting $\bar{a} = \min(k, l) - a$, $\bar{b} = \min(k, m) - b$ and $\bar{c} = \min(l, m) - c$, the left hand side represents the following relation under the combinatorial R:

$$\begin{aligned} x[\zeta_1] \otimes y[\zeta_2] \otimes z[\zeta_3] &\simeq y'[\zeta_2 - \bar{a}] \otimes x'[\zeta_1 + \bar{a}] \otimes z[\zeta_3] \\ &\simeq y'[\zeta_2 - \bar{a}] \otimes z'[\zeta_3 - \bar{b}] \otimes x''[\zeta_1 + \bar{a} + \bar{b}] \\ &\simeq z''[\zeta_3 - \bar{b} - \bar{c}] \otimes y''[\zeta_2 - \bar{a} + \bar{c}] \otimes x''[\zeta_1 + \bar{a} + \bar{b}]. \end{aligned}$$

Similarly, by setting $\bar{d} = \min(l, m) - d$, $\bar{e} = \min(k, m) - e$ and $\bar{f} = \min(k, l) - f$, the same element is transformed along the right hand side as

$$\begin{aligned} x[\zeta_1] \otimes y[\zeta_2] \otimes z[\zeta_3] &\simeq x[\zeta_1] \otimes z^*[\zeta_3 - \bar{d}] \otimes y^*[\zeta_2 + \bar{d}] \\ &\simeq z^{**}[\zeta_3 - \bar{d} - \bar{e}] \otimes x^*[\zeta_1 + \bar{e}] \otimes y^*[\zeta_2 + \bar{d}] \\ &\simeq z^{**}[\zeta_3 - \bar{d} - \bar{e}] \otimes y^{**}[\zeta_2 + \bar{d} - \bar{f}] \otimes x^{**}[\zeta_1 + \bar{e} + \bar{f}]. \end{aligned}$$

Since the Yang-Baxter relation is valid among the affine crystals, we obtain not only $x'' = x^{**}, y'' = y^{**}$ and $z'' = z^{**}$ but also $\bar{b} + \bar{c} = \bar{d} + \bar{e}$, $\bar{a} - \bar{c} = \bar{f} - \bar{d}$ and $\bar{a} + \bar{b} = \bar{e} + \bar{f}$, which are equivalent to the two relations $b + c = d + e$ and $a + b = e + f$. Note that $a + b + c \neq e + f + d$ in general.

Remark 4.3. The energy is invariant under any reordering of $p_1 \otimes \cdots \otimes p_k$ by the combinatorial R. Namely, $\mathcal{E}_i(p_1 \otimes \cdots \otimes p_k) = \mathcal{E}_i(p'_1 \otimes \cdots \otimes p'_k)$ and $\mathcal{E}_i^\vee(p_1 \otimes \cdots \otimes p_k) = \mathcal{E}_i^\vee(p'_1 \otimes \cdots \otimes p'_k)$ hold if $p_1 \otimes \cdots \otimes p_k \simeq p'_1 \otimes \cdots \otimes p'_k$ by the combinatorial R. For $i = n + 1$ this is essentially Proposition 3.9 in [20] and the general i case follows from the symmetry under σ.

Let us consider a particular diagram involving $p_1 \otimes \cdots \otimes p_k$, which is illustrated for $k = 2, 3, 4$. The general case is similar.



Incidentally, this kind of diagrams have been known as the half twist in the construction of link invariants [39].

Lemma 4.4. *The energy $\mathcal{E}_i^\vee(p_1 \otimes \cdots \otimes p_k)$ is the sum of the non-winding numbers Q_i (as e in (4.14)) attached to all the vertices of the corresponding diagram for $p_1 \otimes \cdots \otimes p_k$ as above.*

Proof. For $k = 2$ it is obvious. We use the definition (4.12) and illustrate the induction step along the one from $k = 3$ to $k = 4$.



By the induction assumption, the sum of three $\bullet$ is equal to $\sum_{1 \le j < m \le 3} Q_i(p_j \otimes p_m^{(j+1)})$. Thus we are to verify $e_1 + e_2 + e_3 = \sum_{1 \le j < 4} Q_i(p_j \otimes p_4^{(j+1)})$. But the Yang-Baxter equation

shown above tells that $e_1+e_2+e_3 = d_1+d_2+d_3$, and furthermore, $d_3 = Q_i(p_3 \otimes p_4^{(4)}), d_2 = Q_i(p_2 \otimes p_4^{(3)}), d_1 = Q_i(p_1 \otimes p_4^{(2)})$. □

Lemma 4.5. *$\rho_i(p_1 \otimes \cdots \otimes p_k) - \rho_i(T_\infty(p_1 \otimes \cdots \otimes p_k)) = e_1 + \cdots + e_k$, where e_j's are the i th non-winding numbers specified by the following diagram:*



Proof. In terms of the notation in (4.1), the difference of ρ_i is evaluated as

$$\sum_{j=1}^{k}(x^0_{j,2} + \cdots + x^0_{j,i}) + \sum_{j=1}^{k}(x^1_{j,i+1} + \cdots + x^1_{j,n+1}).$$

By using the graphical rule [17] explained in Appendix B, it is easy to show that the non-winding number Q_i (2.13) is given by $e_j = (x^0_{j,2} + \cdots + x^0_{j,i}) + (x^1_{j,i+1} + \cdots + x^1_{j,n+1})$. □

The main result in this subsection is the following, which identifies the ultradiscrete CTM ρ_i (4.3) with the energy $\mathcal{E}_i$ that originates in the crystal theory.

Proposition 4.6. *$\rho_i(p_1 \otimes \cdots \otimes p_k) = \mathcal{E}_i(p_1 \otimes \cdots \otimes p_k)$ holds for any k and $1 \le i \le n+1$.*

Proof. For $T^t_\infty(p)$ with sufficiently large t, its leftmost k components become $u_{\lambda_1} \otimes \cdots \otimes u_{\lambda_k}$ due to Remark 3.4. In this case the both ρ_i and $\mathcal{E}_i$ are obviously zero. Therefore it suffices to show

$$\rho_i(p_1 \otimes \cdots \otimes p_k) - \rho_i(p'_1 \otimes \cdots \otimes p'_k) = \mathcal{E}_i(p_1 \otimes \cdots \otimes p_k) - \mathcal{E}_i(p'_1 \otimes \cdots \otimes p'_k),$$

where $p'_1 \otimes \cdots \otimes p'_k = T_\infty(p_1 \otimes \cdots \otimes p_k)$. We illustrate the proof for $k = 3$. From Lemma 4.5, we are to show $\mathcal{E}_i(p_1 \otimes p_2 \otimes p_3) = \mathcal{E}_i(p'_1 \otimes p'_2 \otimes p'_3) + e_1 + e_2 + e_3$. Recall that $p'_1 \otimes \cdots \otimes p'_k$ is determined by carrying u_∞ by the combinatorial R through $p_1 \otimes \cdots \otimes p_k$ to the right as $u_\infty \otimes p_1 \otimes \cdots \otimes p_k \simeq p'_1 \otimes \cdots \otimes p'_k \otimes (\cdot)$. Combining this with Lemma 4.4, one can depict the two sides as follows:



$\mathcal{E}_i(p_1 \otimes p_2 \otimes p_3)$



$\mathcal{E}_i(p'_1 \otimes p'_2 \otimes p'_3) + e_1 + e_2 + e_3$

We are to check $a+b+c = a'+b'+c'+e'_1+e'_2+e'_3$. From Remark 4.3, we may assume $\lambda_1 \ge \lambda_2 \ge \lambda_3$ without loss of generality. Then the above equality is a consequence of the separate ones $e'_1 = 0$, $a = a' + e'_2$ and $b + c = b' + c' + e'_3$. To see them, note that $u_\infty \otimes b \simeq u_m \otimes (\cdot)$ for any $b \in B_m$ under the combinatorial R. Moreover $Q_i(u_m \otimes u_j) = 0$

for any m, j. Thus $e_1' = Q_i(u_\infty \otimes u_{\lambda_1}) = 0$ indeed. The other relations can also be seen by appropriately deforming the leftmost line from u_∞ in the right diagram with the aid of the Yang-Baxter equation:



Comparing the lines for p_2 in the left diagram here and the previous one, we find $e_2 + e_2' + a' = e_2 + a + d$. Similarly, the lines for p_3 in the right diagram here and the previous one lead to $e_3 + e_3' + b' + c' = e_3 + b + c + d'$. The proof is finished by noting $d = d' = 0$ because of $Q_i(u_m \otimes u_j) = 0$ for any m, j. □

As a corollary of Proposition 4.6 and (4.4), one has

$$\mathcal{E}_i(u_l \otimes p) = \mathcal{E}_i(p), \tag{4.15}$$

which can also be verified by an argument similar to the above proof.

Remark 4.7. Although the both ρ_i and $\mathcal{E}_i$ admit decompositions into the boundary and the bulk parts as in (4.1) and (4.13), these parts are not equal separately in general. Proposition 4.6 has also been proved by Mark Shimozono by using the technique known as katabolism (private communication).

The energy $\mathcal{E}_{n+1}$ (4.12) and the row transfer matrix energy E_l (3.2) are related by

Proposition 4.8.

$$\mathcal{E}_{n+1}(p) - \mathcal{E}_{n+1}(T_l(p)) = E_l(p).$$

For $l = \infty$ this coincides Lemma 4.5 with $i = n + 1$.

Proof. We illustrate the proof for $p = p_1 \otimes \cdots \otimes p_k$ with $k = 3$. Consider the diagrams:



Here the numbers above the vertices signify the $n + 1$ th non-winding number as in (4.14) with $i = n + 1$, and we have applied the Yang-Baxter relation to the line from

u_l. According to Lemma 4.4, $\mathcal{E}_{n+1}(u_l \otimes p)$ is equal to the sum of all the numbers in the left diagram. Similarly, $\mathcal{E}_{n+1}(T_l(p))$ is obtained from the right diagram as $\mathcal{E}_{n+1}(T_l(p)) = d'_1 + d'_2 + d'_3 + a + b + c$. The Yang-Baxter equation tells that $e_i + d_i = e'_i + d'_i$ for $i = 1, 2, 3$. Using these facts and (4.15), we obtain $\mathcal{E}_{n+1}(p) - \mathcal{E}_{n+1}(T_l(p)) = \mathcal{E}_{n+1}(u_l \otimes p) - \mathcal{E}_{n+1}(T_l(p)) = e'_1 + e'_2 + e'_3$, which coincides with $E_l(p)$ in (3.2). □

4.4. **Proof of Theorem 2.1.** Theorem 2.1 is a simple corollary of (4.5) and

Theorem 4.9. *For any rigged configuration $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ and the corresponding highest state $p = p_1 \otimes \cdots \otimes p_L$ under the KKR bijection, the associated ultradiscrete tau function (2.19) and the ultradiscrete CTM (4.1), (4.3) coincide. Namely*

$$\tau_i(p_1 \otimes \cdots \otimes p_k) = \rho_i(p_1 \otimes \cdots \otimes p_k) \quad (1 \le i \le n+1,\ 1 \le k \le L). \tag{4.16}$$

Proof. Consider the embedding of p into $\mathcal{P}_+(\mu^{(0)}) \otimes B_1^{\otimes L'}$ as $p' = p \otimes 1^{\otimes L'}$. The corresponding rigged configuration is obtained from that of p by just changing $\mu^{(0)}$ into $\mu^{(0)} \sqcup (1^{L'})$. It is easily seen that $\tau_{k,i}$ and $\rho_{k,i}$ for p' are the same as those for p as long as $1 \le k \le L$. Thus we understand them as associated with p' rather than p.

Our proof is based on Propositions 5.1 and 6.1, which will be established in Sections 5 and 6, respectively. Proposition 5.1 states that τ_i satisfies the same bilinear equation (4.9) as ρ_i. Combined with (4.8), it determines $\tau_{k-1,1}, \tau_{k-1,2}, \ldots, \tau_{k-1,n+1}$ successively in this order from $\{\overline{\tau}_{k-1,i}, \overline{\tau}_{k,i}, \tau_{k,i} \mid 1 \le i \le n+1\}$. Namely, the tau function on the NW corner in



is fixed from those on the NE, SW and SE. Like ρ and $\overline{\rho}$, the tau functions τ and $\overline{\tau}$ are associated with p' and $T_\infty(p')$, respectively (see the beginning of Section 5), and the above diagram can be extended to a two-dimensional square lattice with the indicated coordinates. The square at (k, t) is associated with $\big(\tau_{k,i}(T^t_\infty(p'))\big)_{i=1}^{n+1}$.

Consider the rectangular region on the lattice $0 \le t \le t_0, 1 \le k \le L + L'$, where the tau functions for p' constitutes the top line $t = 0$ of it. They are uniquely determined from the right boundary $k = L + L'$, i.e., $\{\big(\tau_{L+L',i}(T^t_\infty(p'))\big)_{i=1}^{n+1} \mid 0 \le t \le t_0\}$, and the bottom boundary $t = t_0$, i.e., $\{\big(\tau_{k,i}(T^{t_0}_\infty(p'))\big)_{i=1}^{n+1} \mid 1 \le k \le L + L'\}$. The coincidence of ρ_i and τ_i on these boundaries will be proved in Proposition 6.1 by taking t_0 and L' sufficiently large. □

5. Bilinear relation for τ_i

Let $\tau_{k,d}$ be the ultradiscrete tau function specified in Theorem 2.1 and (2.18)–(2.20). We define $\overline{\tau}_{k,d}$ to be $\tau_{k,d}$ with $|s^{(1)}|$ replaced by $|s^{(1)}| + |\nu^{(1)}|$ in (2.20). In view of Proposition 3.5, this corresponds to the rigged configuration that has undergone the time evolution T_∞ once.

Proposition 5.1. *The substitution $\rho_{k,d} = \tau_{k,d}$ and $\overline{\rho}_{k,d} = \overline{\tau}_{k,d}$ solves the bilinear equation (4.9).*

This section is devoted to the proof of Proposition 5.1 by a refinement of the approach in [26]. We invoke the free fermion construction of tau functions associated with $\mathfrak{gl}(\infty)$ [37]. For $l \in \mathbb{Z}$, set

$$\sigma_l(x) = \langle l | e^{H(x)} g | l \rangle, \quad g = \exp\Big(\sum_{(a,i)} c_i^{(a)} \psi(p_i^{(a)}) \psi^*(q_i^{(a)})\Big), \tag{5.1}$$

where the notation is the same as eq.(2.3) in [37] except that τ_l there is denoted by σ_l here for distinction from (2.19). ($p_i^{(a)}$ here is not the vacancy number (2.7).) The operators $\psi(k) = \sum_{j\in\mathbb{Z}} \psi_j k^j, \psi^*(k) = \sum_{j\in\mathbb{Z}} \psi_j^* k^{-j}$ are the free fermions. They obey the anti-commutation relations $[\psi_i, \psi_j]_+ = [\psi_i^*, \psi_j^*]_+ = 0$ and $[\psi_i, \psi_j^*]_+ = \delta_{ij}$, hence $\psi(k)^2 = \psi^*(k)^2 = 0$. $|l\rangle$ is the charge l vacuum of the Fock space. $H(x) = \sum_{i\geq 1} x_i \sum_{j\in\mathbb{Z}} \psi_j \psi_{j+i}^*$ is the Hamiltonian with infinitely many time variables $x = (x_1, x_2, \ldots)$. In (5.1), we associate each triple $(c_i^{(a)}, p_i^{(a)}, q_i^{(a)})$ with the data $(\mu_i^{(a)}, r_i^{(a)})$ in the rigged configuration $(\lambda, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$. The sum extends over all the colors $1 \leq a \leq n$ and the rows $1 \leq i \leq \ell(\mu^{(a)})$. The tau function (5.1) is an N-soliton solution of the KP hierarchy with $N = \ell(\mu^{(1)}) + \cdots + \ell(\mu^{(n)})$.

The time evolution of the free fermion is given by $e^{H(x)}\psi(k)e^{-H(x)} = e^{\xi(x,k)}\psi(k)$ and $e^{H(x)}\psi^*(k)e^{-H(x)} = e^{-\xi(x,k)}\psi^*(k)$ with $\xi(x,k) = \sum_{i\geq 1} x_i k^i$. Consequently,

$$e^{H(x)}\psi(p)\psi^*(q)e^{-H(x)} = \frac{\beta - q}{\beta - p}\psi(p)\psi^*(q)$$

for $x = \varepsilon(\beta^{-1}) := (\beta^{-1}, \frac{1}{2}\beta^{-2}, \frac{1}{3}\beta^{-3}, \ldots)$. For $z_k := \varepsilon(\beta_1^{-1}) + \cdots + \varepsilon(\beta_k^{-1})$, the tau function is expanded as

$$\sigma_l(z_k) = \sum_{\nu=(\nu^{(1)},\ldots,\nu^{(n)})} \sigma_l(z_k)_\nu, \tag{5.2}$$

$$\sigma_l(z_k)_\nu = \Delta_\nu \prod_{(a,i)} c_i^{(a)} q_i^{(a)} \Big(\frac{p_i^{(a)}}{q_i^{(a)}}\Big)^l \prod_{j=1}^{k} \frac{\beta_j - q_i^{(a)}}{\beta_j - p_i^{(a)}}, \tag{5.3}$$

$$\Delta_\nu = \frac{\prod_{(a,i)<(b,j)} (p_i^{(a)} - p_j^{(b)})(q_j^{(b)} - q_i^{(a)})}{\prod_{(a,i),(b,j)} (p_i^{(a)} - q_j^{(b)})}, \tag{5.4}$$

where the sum (5.2) extends over the subsets $\nu^{(1)} \subseteq \mu^{(1)}, \ldots, \nu^{(n)} \subseteq \mu^{(n)}$ independently. In (5.3), the product $\prod_{(a,i)}$ runs over the rows of the selected subset $\nu^{(a)} \subseteq \mu^{(a)}$. In (5.4), $\prod_{(a,i)<(b,j)}$ runs over the pairs of such indices, whereas $\prod_{(a,i),(b,j)}$ simply means the double product. Δ_ν is the Cauchy determinant of the free fermion up to an overall power of $p_i^{(a)}$ and $q_i^{(a)}$. It is derived by using the formulas:

$$\langle l|\psi(p)\psi^*(q)|l\rangle = \sum_{j\leq l-1} p^j q^{-j} = \frac{q}{p-q}\left(\frac{p}{q}\right)^l,$$

$$\langle l|\psi(p_1)\cdots\psi(p_m)\psi^*(q_m)\cdots\psi^*(q_1)|l\rangle = \frac{\prod_{i<j}(p_i - p_j)(q_j - q_i)}{\prod_{i,j=1}^m (p_i - q_j)} \prod_{i=1}^m q_i \left(\frac{p_i}{q_i}\right)^l.$$

Now we make a special choice of the parameters that further reflects the rigged configuration $(\lambda, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$. Fixing $d \in \{2, \ldots, n+1\}$ we set

$$p_i^{(a)} = \kappa^{(a)} - \delta_i^{(a)} \exp(-\frac{\mu_i^{(a)}}{\epsilon}), \quad q_i^{(a)} = \kappa^{(a+1)} + \delta_i^{(a)} \exp(-\frac{\mu_i^{(a)}}{\epsilon}), \tag{5.5}$$

$$c_i^{(a)} q_i^{(a)} = \begin{cases} \delta_i'^{(a)} \exp(-\frac{2\mu_i^{(a)} + r_i^{(a)}}{\epsilon}) & \text{if } a \in \{1, d\}, \\ \delta_i'^{(a)} \exp(-\frac{\mu_i^{(a)} + r_i^{(a)}}{\epsilon}) & \text{otherwise,} \end{cases} \tag{5.6}$$

$$\beta_j = \kappa^{(1)} + \delta_j'' \exp(-\frac{\lambda_j}{\epsilon}), \tag{5.7}$$

where $1 \le a \le n$ and $\epsilon > 0$. Here $\kappa^{(1)}, \dots, \kappa^{(n+1)}$ and $\delta_i^{(a)}, \delta_i^{'(a)}, \delta_j''$ are ϵ-independent generic (hence distinct) parameters such that

$$\kappa^{(1)} > \cdots > \kappa^{(d-1)} > \kappa^{(d)} = 0 > \kappa^{(d+1)} > \cdots > \kappa^{(n+1)}, \tag{5.8}$$

$$\delta_i^{(a)} > 0,\ \delta_i^{'(a)} > 0,\ \delta_j'' > 0. \tag{5.9}$$

Lemma 5.2. *Set $q = e^{-1/\epsilon}$. In the limit $q \to 0$, the summand $\sigma_l(z_k)_\nu$ (5.3) of the tau function has the following behavior:*

$$\begin{aligned} \sigma_l(z_k)_\nu &= q^{c(\nu,s)+|\nu^{(1)}|+|\nu^{(d)}|-l(|\nu^{(d-1)}|-|\nu^{(d)}|)}(\chi_\nu + \mathcal{O}(q)), \quad \chi_\nu > 0, \\ \sigma_l(z_k + \varepsilon(\kappa^{(1)-1}))_\nu &= q^{c(\nu,s)+|\nu^{(d)}|-l(|\nu^{(d-1)}|-|\nu^{(d)}|)}(\chi'_\nu + \mathcal{O}(q)), \quad \chi'_\nu > 0, \end{aligned} \tag{5.10}$$

where χ_ν and χ'_ν are independent of ϵ. $c(\nu, s)$ is defined by (2.20) with $\lambda = (\lambda_1, \dots, \lambda_k)$.

We denote by $A \xrightarrow{\mathrm{UD}} a$ the relation $a = \lim_{\epsilon \to +0} \epsilon \log A$ under the ultradiscretization. It means that $A = A_0 q^{-a} +$ higher order terms in q for some leading coefficient $A_0 (\neq 0)$. (A_0 still can depend on ϵ as long as $A_0 \xrightarrow{\mathrm{UD}} 0$ although it is not needed in our case.) We let the relation $A \sim B$ mean $\lim_{\epsilon \to +0} \epsilon \log A = \lim_{\epsilon \to +0} \epsilon \log B$.

Proof. Let $\{(\nu_i^{(a)}, s_i^{(a)})\}$ be the subset of the rigged configuration $\{(\mu_i^{(a)}, r_i^{(a)})\}$ corresponding to $\nu = (\nu^{(1)}, \dots, \nu^{(n)})$ as in (2.20). We investigate the leading power of the constituent factors in (5.3). From (5.5)–(5.7) we find

$$\text{(i)} \prod_{(a,i)} c_i^{(a)} q_i^{(a)} \xrightarrow{\mathrm{UD}} -\sum_{a=1}^{n} (|\nu^{(a)}| + |s^{(a)}|) - |\nu^{(1)}| - |\nu^{(d)}|,$$

$$\text{(ii)} \prod_{(a,i)} \left(\frac{p_i^{(a)}}{q_i^{(a)}} \right)^l \sim \prod_i \left(\frac{p_i^{(d)}}{q_i^{(d-1)}} \right)^l \xrightarrow{\mathrm{UD}} l(|\nu^{(d-1)}| - |\nu^{(d)}|),$$

$$\text{(iii)} \prod_{(a,i)} \prod_{j=1}^{k} \frac{\beta_j - q_i^{(a)}}{\beta_j - p_i^{(a)}} \sim \prod_i \prod_{j=1}^{k} \frac{1}{\beta_j - p_i^{(1)}} \xrightarrow{\mathrm{UD}} \min(\lambda, \nu^{(1)}),$$

$$\begin{aligned} \text{(iv)} \prod_{(a,i)<(b,j)} (p_i^{(a)} - p_j^{(b)})(q_j^{(b)} - q_i^{(a)}) &\sim \prod_{a=1}^{n} \prod_{i<j} (p_i^{(a)} - p_j^{(a)})^2 \\ &\xrightarrow{\mathrm{UD}} \sum_{a=1}^{n} (-\min(\nu^{(a)}, \nu^{(a)}) + |\nu^{(a)}|), \end{aligned}$$

$$\text{(v)} \prod_{(a,i),(b,j)} (p_i^{(a)} - q_j^{(b)})^{-1} \sim \prod_{a=2}^{n} \prod_{i,j} (p_i^{(a)} - q_j^{(a-1)})^{-1} \xrightarrow{\mathrm{UD}} \sum_{a=2}^{n} \min(\nu^{(a)}, \nu^{(a-1)}),$$

$$\text{(vi)} \prod_{(a,i)} \frac{\kappa^{(1)} - q_i^{(a)}}{\kappa^{(1)} - p_i^{(a)}} \sim \prod_i \frac{1}{\kappa^{(1)} - p_i^{(1)}} \xrightarrow{\mathrm{UD}} |\nu^{(1)}|,$$

where $|\nu^{(n+1)}| = 0$ and the notation (2.3) is used. The contributions (i)–(v) sum up to $-c(\nu, s) - |\nu^{(1)}| - |\nu^{(d)}| + l(|\nu^{(d-1)}| - |\nu^{(d)}|)$. This verifies the leading power of $\sigma_l(z_k)_\nu$ in (5.10). Similarly, the one for $\sigma_l(z_k + \varepsilon(\kappa^{(1)-1}))_\nu$ is derived by including the contribution from (vi).

The remaining task is to check the positivity and ϵ-independence of the leading coefficients χ_ν and χ'_ν. We first illustrate them along $\sigma_l(z_k)_\nu$. In the right hand side of (5.3), we show the positivity individually for the constituent factors (i), (ii), (iii) and $\Delta_\nu =$(iv)$\times$(v) considered in the above. The leading coefficient from (i) is $\prod_{(a,i)} \delta_i^{'(a)}$ by

(5.6), which is positive due to (5.9). The leading coefficient from (ii) is 1 if $l = 0$. If $l = -1$, it is given by

$$\prod_i \frac{\delta_i^{(d-1)}}{\kappa^{(d-1)}} \prod_j \frac{\kappa^{(d+1)}}{-\delta_j^{(d)}} \prod_{\substack{1\le a\le n\\ a\neq d-1,d}} \left(\frac{\kappa^{(a+1)}}{\kappa^{(a)}}\right)^{\ell(\nu^{(a)})},$$

where the products on i and j extend over the selected rows in $\nu^{(d-1)}$ and $\nu^{(d)}$, respectively. The symbol $\ell(\nu^{(a)})$ denotes the length of $\nu^{(a)}$ as defined in (2.1). This is positive thanks to (5.8) and (5.9). The leading coefficient from (iii) with a fixed j is equal to the one from

$$\prod_i \frac{\kappa^{(1)} - \kappa^{(2)}}{\delta_j'' q^{\lambda_j} + \delta_i^{(1)} q^{\mu_i^{(1)}}} \prod_{\substack{(a,i)\\ a\ge 2}} \frac{\kappa^{(1)} - \kappa^{(a+1)}}{\kappa^{(1)} - \kappa^{(a)}}.$$

It is positive by (5.8) and (5.9). The leading coefficients from (iv) and (v) are respectively equal to those in

$$\prod_{a=1}^n \prod_{i<j} (\delta_j^{(a)} q^{\nu_j^{(a)}} - \delta_i^{(a)} q^{\nu_i^{(a)}})^2 \prod_{1\le a<b\le n} \big((\kappa^{(a)} - \kappa^{(b)})(\kappa^{(b+1)} - \kappa^{(a+1)})\big)^{\ell(\nu^{(a)})\ell(\nu^{(b)})},$$

$$\prod_{a=2}^n \prod_{i,j} (-\delta_i^{(a)} q^{\nu_i^{(a)}} - \delta_j^{(a-1)} q^{\nu_j^{(a-1)}})^{-1} \prod_{\substack{1\le a,b\le n\\ a\neq b+1}} (\kappa^{(a)} - \kappa^{(b+1)})^{-\ell(\nu^{(a)})\ell(\nu^{(b)})}.$$

In view of (5.8) and (5.9), the coefficients are both positive apart from the same sign factor $(-1)^{\sum_{1\le a<b\le n} \ell(\nu^{(a)})\ell(\nu^{(b)})}$. Thus the leading coefficient from the product (iv)×(v) is positive.

For $\sigma_l(z_k + \varepsilon(\kappa^{(1)-1}))$, the leading positivity $\chi'_\nu > 0$ is proved similarly. The only necessary modification is to include the contribution from (vi):

$$\frac{\prod_{a=1}^n (\kappa^{(1)} - \kappa^{(a+1)})^{\ell(\nu^{(a)})}}{\prod_i \delta_i^{(1)} \prod_{a=2}^n (\kappa^{(1)} - \kappa^{(a)})^{\ell(\nu^{(a)})}},$$

which is again positive due to (5.8) and (5.9). Finally, χ_ν and χ'_ν are ϵ-independent as they are rational functions of the parameters appearing in (5.8) and (5.9) only. □

Lemma 5.3.

$$\sigma_0(z_k) \xrightarrow{\text{UD}} \overline{\tau}_{k,d}, \qquad \sigma_{-1}(z_k) \xrightarrow{\text{UD}} \overline{\tau}_{k,d-1},$$
$$\sigma_0(z_k + \varepsilon(\kappa^{(1)-1})) \xrightarrow{\text{UD}} \tau_{k,d}, \quad \sigma_{-1}(z_k + \varepsilon(\kappa^{(1)-1})) \xrightarrow{\text{UD}} \tau_{k,d-1}.$$

Proof. For example we consider the UD limit

$$\lim_{\epsilon\to+0} \epsilon \log \sigma_0(z_k + \varepsilon(\kappa^{(1)-1})) = \lim_{\epsilon\to+0} \epsilon \log\Big(\sum_\nu \chi'_\nu\, q^{c(\nu,s)+|\nu^{(d)}|}\Big), \tag{5.11}$$

where $q = e^{-1/\epsilon}$ and (5.10) has been substituted. Lemma 5.2 furthermore tells that there is no cancellation in the ν-sum here because of $\chi'_\nu > 0$. Therefore the limit tends to $\max_\nu\{-c(\nu,s) - |\nu^{(d)}|\} = \tau_{k,d}$. See the definitions of $\tau_k(\lambda)$ (2.19) and $\tau_{k,d}$ (2.21). The other limits are confirmed similarly. □

Proof of Proposition 5.1. It is well known that σ_l satisfies the bilinear equation:

$$\begin{aligned}
&(\alpha^{-1} - \beta^{-1})\sigma_0(z + \varepsilon(\alpha^{-1}) + \varepsilon(\beta^{-1}))\sigma_{-1}(z + \varepsilon(\gamma^{-1}))\\
&+(\beta^{-1} - \gamma^{-1})\sigma_0(z + \varepsilon(\beta^{-1}) + \varepsilon(\gamma^{-1}))\sigma_{-1}(z + \varepsilon(\alpha^{-1}))\\
&+(\gamma^{-1} - \alpha^{-1})\sigma_0(z + \varepsilon(\gamma^{-1}) + \varepsilon(\alpha^{-1}))\sigma_{-1}(z + \varepsilon(\beta^{-1})) = 0.
\end{aligned}$$

This is derived by setting $x = z + \varepsilon(\alpha^{-1}) + \varepsilon(\beta^{-1}) + \varepsilon(\gamma^{-1})$, $x' = z$ and $(l, l') = (0, -1)$ in eq.$(2.4)_{l,l'}$ in p956 of [37]. Setting

$$\alpha = \kappa^{(1)}, \ \ \beta = \beta_k, \ \ \gamma = \infty, \ \ x = z_{k-1} = \varepsilon(\beta_1^{-1}) + \cdots + \varepsilon(\beta_{k-1}^{-1}),$$

we get

$$\begin{aligned}
&\beta_k \sigma_0(z_{k-1} + \varepsilon(\kappa^{(1)-1}))\sigma_{-1}(z_k) \\
&= \kappa^{(1)} \sigma_0(z_k)\sigma_{-1}(z_{k-1} + \varepsilon(\kappa^{(1)-1})) + \delta_k'' e^{-\lambda_k/\epsilon} \sigma_0(z_k + \varepsilon(\kappa^{(1)-1}))\sigma_{-1}(z_{k-1}),
\end{aligned}$$

where $\beta_k - \kappa^{(1)}$ has been evaluated by (5.7). In view of (5.8) and (5.9), the coefficients $\beta_k, \kappa^{(1)}$ and δ_k'' here are all positive and ϵ-independent. Moreover from Lemma 5.2, there is no cancellation of the leading terms coming from the two terms on the right hand side. Therefore by taking the UD limit $\lim_{\epsilon \to +0} \epsilon \log(\cdot)$ of the two sides and applying Lemma 5.3, we obtain

$$\tau_{k-1,d} + \overline{\tau}_{k,d-1} = \max(\overline{\tau}_{k,d} + \tau_{k-1,d-1},\ \tau_{k,d} + \overline{\tau}_{k-1,d-1} - \lambda_k). \tag{5.12}$$

This coincides with (4.9) with ρ replaced by τ. Note that the range $2 \le d \le n+1$ for the both also match. This completes the proof of Proposition 5.1. □

Let us compare the results in this section with the similar ones in section IV of [26]. In [26], the tau function is supposed to fulfill the periodicity $\tau_{k,d} = \tau_{k,d+n+1}$ in the present notation. This led to a reduction condition (Prop.4.4 in [HHIKTT]) on each pair of the parameters $(p_i^{(a)}, q_i^{(a)})$ in (5.1), restricting the class of tau functions captured in the UD limit. In our approach, reduction conditions are bypassed by the special choice of the parameters (5.5)–(5.9) depending on the d that enters the bilinear equation (5.12) to prove. As it will turn out in Section 7.3, the ultradiscrete tau functions derived here cover *all* the solutions of the box-ball system.

6. Asymptotic coincidence of τ_i and ρ_i

6.1. **Statement and its reduction.**

In this section we prove

Proposition 6.1. *Given a highest path p with length L, set $p' = p \otimes \overbrace{1 \otimes \cdots \otimes 1}^{L'}$ and $k_0 = L + L'$. Then the equalities* $(1 \le i \le n+1)$

$$\tau_{k,i}(T_\infty^{t_0}(p')) = \rho_{k,i}(T_\infty^{t_0}(p')) \quad 1 \le k \le k_0, \tag{6.1}$$

$$\tau_{k_0,i}(T_\infty^{t}(p')) = \rho_{k_0,i}(T_\infty^{t}(p')) \quad 0 \le t \le t_0 \tag{6.2}$$

hold if $t_0 \gg 1$ in (6.1), and if furthermore $k_0 \gg L t_0$ in (6.2).

Combined with Proposition 5.1, it establishes Theorem 4.9 and thereby completes the proof of Theorem 2.1. Let $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ be the rigged configuration corresponding to $T_\infty^{t_0}(p')$. Without loss of generality we assume $\mu_1^{(1)} \le \mu_2^{(1)} \le \cdots$. Moreover from the condition $t_0 \gg 1$ and Proposition 3.5, we assume

$$\begin{aligned}
&1 \ll r_1^{(1)} \le r_2^{(1)} \le r_3^{(1)} \le \cdots, \\
&r_i^{(1)} \ll r_j^{(1)} \quad \text{if} \ \ \mu_i^{(1)} < \mu_j^{(1)}
\end{aligned} \tag{6.3}$$

throughout this section. From Remark 3.4 and $k_0 \gg L t_0$, the state $T_\infty^{t_0}(p')$ takes the form:

$$T_\infty^{t_0}(p') = u_{\lambda_1} \otimes \cdots \otimes u_{\lambda_L} \otimes \underbrace{\overbrace{1 \otimes \cdots \otimes 1}^{a \gg 1} \otimes (\cdots\cdots\cdots) \otimes \overbrace{1 \otimes \cdots \otimes 1}^{b \gg 1}}_{\tilde{p}}, \tag{6.4}$$

where $\lambda = (\lambda_1, \ldots, \lambda_L)$ are the numbers such that $p \in B_{\lambda_1} \otimes \cdots \otimes B_{\lambda_L}$ and $\tilde{p} \in B_1^{\otimes L'}$ is a highest path.

Lemma 6.2. *Under the same condition as Proposition 6.1, the following relation holds:*

$$\tau_{k_0,i}(T^t_\infty(p')) - \tau_{k_0,i}(T^{t+1}_\infty(p')) = \rho_{k_0,i}(T^t_\infty(p')) - \rho_{k_0,i}(T^{t+1}_\infty(p')) \quad (0 \le t \le t_0 - 1).$$

Proof. Suppose $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \dots, (\mu^{(n)}, r^{(n)}))$ is the rigged configuration for $T^{t_0}_\infty(p')$. From the definition (4.1) and the assumed situation (6.4), it is easily seen that $\rho_{k_0,i}(T^t_\infty(p')) - \rho_{k_0,i}(T^{t+1}_\infty(p'))$ is the number of balls with colors $2, \dots, n+1$ contained in p, which is equal to $|\mu^{(1)}|$. To calculate $\tau_{k_0,i}(T^t_\infty(p')) - \tau_{k_0,i}(T^{t+1}_\infty(p'))$, we apply the formula (2.23). $\tau_{k_0,i}(T^t_\infty(p'))$ is obtained by replacing λ there with $\lambda \sqcup (1^{k_0-L})$ and $r^{(1)}_i$ with $r^{(1)}_i - (t_0 - t)\mu^{(1)}_i$ by Proposition 3.5. Then the max contains k_0 only via $\min(\lambda \sqcup (1^{k_0-L}), \nu)$, hence one can let it be achieved at $\nu = \mu^{(1)}$ by taking k_0 sufficiently large. Consequently $\tau_{k_0,i}(T^t_\infty(p')) = \min(\lambda \sqcup (1^{k_0-L}), \mu^{(1)}) - \min(\mu^{(1)}, \mu^{(1)}) - |r^{(1)}| + (t_0 - t)|\mu^{(1)}| + \tau^{(1)}_i(\mu^{(1)})$ for any $0 \le t \le t_0$ as long as $k_0 \gg Lt_0$. Therefore $\tau_{k_0,i}(T^t_\infty(p')) - \tau_{k_0,i}(T^{t+1}_\infty(p')) = |\mu^{(1)}|$ in agreement with $\rho_{k_0,i}(T^t_\infty(p')) - \rho_{k_0,i}(T^{t+1}_\infty(p'))$. □

By Lemma 6.2, (6.2) is attributed to $t = t_0$ case. Thus the proof of Proposition 6.1 reduces to showing (6.1), on which we shall concentrate from now on.

Lemma 6.3. *For $1 \le k \le L$, $\rho_{k,i}(T^{t_0}_\infty(p')) = \tau_{k,i}(T^{t_0}_\infty(p')) = 0$. For $L < k \le k_0$, the following relations hold:*

$$\rho_{k,i}(T^{t_0}_\infty(p')) = \rho_{k-L,i}(\tilde{p}), \tag{6.5}$$

$$\tau_{k,i}(T^{t_0}_\infty(p')) = \tau_{k-L,i}(\tilde{p}), \tag{6.6}$$

where $\tilde{p} \in B_1^{\otimes L'}$ is defined in (6.4).

Proof. For $\rho_{k,i}$, the assertion is obvious from (6.4) and the definition (4.1). As for $\tau_{k,i}$, we use the expression (2.23) for $T^{t_0}_\infty(p')$ which corresponds to the rigged configuration $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \dots, (\mu^{(n)}, r^{(n)}))$.

$$\tau_i(\xi) = \max_{\nu \subseteq \mu^{(1)}} \{\min(\xi, \nu) - \min(\nu, \nu) - |s| + \tau^{(1)}_i(\nu)\} \quad \xi \subseteq \mu^{(0)}. \tag{6.7}$$

According to (6.4), we have $\mu^{(0)} = \lambda \sqcup (1^{L'})$. From Proposition 3.5, we know that $r^{(1)}_i = \mu^{(1)}_i t_0 + \bar{r}^{(1)}_i$, where $\bar{r}^{(1)}_i$ is the rigging for p'. (This is also equal to the rigging for p in Proposition 6.1 although this fact is not used below.) Thus t_0 enters (6.7) only via $|s| = |\nu| t_0 + |\bar{s}|$, where $|\bar{s}|$ is t_0-independent. Fixing $\xi = (\lambda_1, \dots, \lambda_k)$ with $1 \le k \le L$ and taking t_0 sufficiently large, we see that the maximum (6.7) forces the choice $\nu = \emptyset$. This yields $\tau_{k,i}(T^{t_0}_\infty(p')) = \tau^{(1)}_i(\emptyset) = 0$ for $1 \le k \le L$, where the latter equality is due to Lemma 2.3.

The maximum can be different from 0 for $L < k \le k_0$, where we are allowed to take k so large up to k_0 depending on t_0. This corresponds to the situation (6.6), which will be considered in the sequel. To compute the right hand side of (6.6) by (6.7), we need to know the rigged configuration for $\tilde{p}$. In view of (6.4), it is obtained from the one $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \dots, (\mu^{(n)}, r^{(n)}))$ for $T^{t_0}_\infty(p')$ by replacing $\mu^{(0)}$ with $(1^{L'})$ and the rigging $r^{(1)}_i$ with $\tilde{r}^{(1)}_i = r^{(1)}_i - \sum_{j=1}^L \min(\lambda_j, \mu^{(1)}_i)$. See Lemma C.3. This amounts to changing $|s|$ in (6.7) to $|s| - \min(\lambda, \nu)$ in the notation (2.3). Thus by setting $\xi = (1^{k-L})$, we get

$$\begin{aligned}\tau_{k-L,i}(\tilde{p}) &= \max_{\nu \subseteq \mu^{(1)}} \{\min((1^{k-L}), \nu) - \min(\nu, \nu) - (|s| - \min(\lambda, \nu)) + \tau^{(1)}_i(\nu)\} \\ &= \max_{\nu \subseteq \mu^{(1)}} \{\min(\lambda \sqcup (1^{k-L}), \nu) - \min(\nu, \nu) - |s| + \tau^{(1)}_i(\nu)\}.\end{aligned}$$

Since $\lambda \sqcup (1^{k-L}) = (\mu^{(0)}_1, \dots, \mu^{(0)}_k)$, this is nothing but the expression of $\tau_{k,i}(T^{t_0}_\infty(p'))$ by (6.7). □

Thanks to Lemma 6.3, we may assume $\lambda = \emptyset$ in (6.4) without loss of generality.

To summarize so far, we have reduced Proposition 6.1 to (6.1) for p' such that $p' \in B_1^{\otimes k_0}$. Resetting the meaning of $p, p', \tilde{p}, L, L'$ and k_0, we restate it as

Proposition 6.4. *Let $p \in B_1^{\otimes L}$ be a highest path and $((1^L), (\mu^{(1)}, r^{(1)}), \dots, (\mu^{(n)}, r^{(n)}))$ be its rigged configuration with $\mu_1^{(1)} \le \mu_2^{(1)} \le \cdots$. If L is sufficiently large and the condition (6.3) is satisfied, the equality*

$$\tau_{k,i}(p) = \rho_{k,i}(p) \tag{6.8}$$

is valid for $1 \le k \le L$, $2 \le i \le n+1$.

A highest path $p \in B_1^{\otimes L}$ satisfying the assumption of Proposition 6.4 will be called an *asymptotic state*. We have excluded $i = 1$ case since it is contained as the $i = n+1$ case of $T_\infty(p)$ which is also an asymptotic state. See (2.19), Proposition 3.5 and (4.2). The remainder of this section is devoted to the proof of Proposition 6.4. Our strategy is to express the both sides of (6.8) in terms of the quantities associated with the smaller algebra $A_{n-1}^{(1)}$ and invoke the induction with respect to n. Note that the induction allows us to use Theorem 2.2 with $1 \le a \le n-1$ and Theorem 4.9 for $A_{n-1}^{(1)}$.

6.2. **Precise description of asymptotic states.** The KKR bijection from rigged configurations to highest paths is known to be equivalent with the vertex operator construction [34, 35]. Here we utilize the notions in the latter formalism such as scattering data and normal ordering explained in Appendix D. In particular, we remark that a scattering data $b_1[d_1] \otimes \cdots \otimes b_N[d_N] \in \mathrm{Aff}(B_{\mu_1}^{\ge 2}) \otimes \cdots \otimes \mathrm{Aff}(B_{\mu_N}^{\ge 2})$ for an asymptotic state is normal ordered if and only if $\mu_1 \le \cdots \le \mu_N$.

Lemma 6.5. *For an asymptotic state p, denote any successive tensor product components of the normal ordered scattering data by*

$$\cdots \otimes \boxed{B}_{d_2} \otimes \boxed{A}_{d_1} \otimes \cdots . \tag{6.9}$$

Let the semistandard tableaux A and B be

$$\boxed{A} = \boxed{a_1 a_2 \dots a_{l_A}} \in B_{l_A}^{\ge 2}, \qquad 2 \le a_1 \le a_2 \le \cdots \le a_{l_A} \le n+1,$$
$$\boxed{B} = \boxed{b_1 b_2 \dots b_{l_B}} \in B_{l_B}^{\ge 2}, \qquad 2 \le b_1 \le b_2 \le \cdots \le b_{l_B} \le n+1\,.$$

Then locally p has the form:

$$\cdots 11\, b_{l_B} \cdots b_2 b_1 \overbrace{11\cdots 1}^{d_1-d_2} a_{l_A} \cdots a_2 a_1 11 \cdots . \tag{6.10}$$

Proof. Since p is an asymptotic state, we have $l_B \le l_A$. We divide the proof into two cases.

Case 1. Assume $l_B < l_A$. From the definition of the modes of scattering data (D.3), we have $l_B \ll d_1 - d_2$ for asymptotic states. Therefore, the calculation of the vertex operator goes as (see around (E.2) for the explanation of Φ_B)

$$\begin{aligned}
&\Phi_B(\overbrace{11\cdots 1}^{d_1-d_2} a_{l_A} \cdots a_2 a_1 \cdots) \\
&= b_{l_B} \cdots b_2 b_1 \overbrace{11\cdots 1}^{d_1-d_2-l_B} T_{l_B}(a_{l_A} \cdots a_2 a_1 \cdots) \\
&= b_{l_B} \cdots b_2 b_1 \overbrace{11\cdots 1}^{d_1-d_2} a_{l_A} \cdots a_2 a_1 \cdots ,
\end{aligned}$$

where T_l is a time evolution of the box-ball system with capacity l career (3.1).

Case 2. Next, consider the case $l_B = l_A = l$. Let the energy function be $H = H(B \otimes A)$. Applying the definition of the mode (D.3) to (6.9), we have $d_2 = l + r_B + h$ and $d_1 =$

$l + r_A + h + H(B \otimes A)$, where r_A, r_B are the riggings for A, B and h denotes the last term in (D.3) for d_2 here. Since the asymptotic state satisfies the condition (6.3), we have $r_B \le r_A$, leading to $H \le d_1 - d_2 (=: \Delta)$. If $\Delta \ge l$, the proof is the same as *Case 1*. Therefore assume $H \le \Delta < l$ in the following. Calculating the action of Φ_B, we arrive at the following situation:



where $B' = \boxed{1^{\Delta} b_1 b_2 \cdots b_{l-\Delta}}$. The diagram says that $B' \otimes a_l \simeq a' \otimes (\cdots)$ under the combinatorial R. Let us show that $a' = b_{l-\Delta}$. For the purpose, we first claim $b_{l-\Delta} < a_l$. In fact, suppose $b_{l-\Delta} \ge a_l$ on the contrary. We construct the pairs for $B \otimes A$ according to the graphical rule in Appendix B to compute $H = H(B \otimes A)$. We know that there are H winding pairs irrespective of the ways of making pairs. Since b_i is weakly increasing with respect to i, we see that more than $\Delta + 1 (> H)$ i's satisfy $b_i \ge a_l$. On the other hand, a_l is the largest letter in A, therefore all the letters in B greater than a_l have to constitute winding pairs, and we have seen that the number of these winding pairs is greater than H. This is a contradiction. Therefore we obtain $b_{l-\Delta} < a_l$.

We have seen that $b_{l-\Delta} < a_l$, and we know that $b_{l-\Delta}$ is the largest number in B'. When we construct the pairs for $B' \otimes a_l$, this fact means that $b_{l-\Delta}$ and a_l form an unwinding pair. Therefore, the action of the combinatorial R is given by $B' \otimes a_l \simeq a' \otimes (\cdots)$ with $a' = b_{l-\Delta}$. By continuing the same argument, we arrive at (6.10). □

In what follows we use the notation explained in Section 2.1.

Lemma 6.6. *Suppose that Proposition 6.4 is true for $A^{(1)}_{n-1}$. For a rigged configuration $((1^L), (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ with $\mu^{(1)} = (\mu_1, \ldots, \mu_N)$ and $r^{(1)} = (r_1, \ldots, r_N)$, assume $\mu_1 \le \cdots \le \mu_N$. Then the corresponding scattering data $b_1[d_1] \otimes \cdots \otimes b_N[d_N] \in \mathit{Aff}(B^{\ge 2}_{\mu_1}) \otimes \cdots \otimes \mathit{Aff}(B^{\ge 2}_{\mu_N})$ is given by*

$$b_M = (x_2, \ldots, x_{n+1}), \quad x_i = \tau^{(1)}_{M,i} - \tau^{(1)}_{M-1,i} - \tau^{(1)}_{M,i-1} + \tau^{(1)}_{M-1,i-1}, \tag{6.11}$$

$$d_M = |\mu_{[M]}| + r_M + \tau^{(1)}_{M-1,n+1} - \tau^{(1)}_{M,n+1}. \tag{6.12}$$

This lemma is shown without assuming that the scattering data $b_1[d_1] \otimes \cdots \otimes b_N[d_N]$ is normal ordered.

Proof. From the arguments in Section 6.1, the assumption makes Theorem 2.1 for $A^{(1)}_{n-1}$ valid. Then (6.11) is a corollary of Theorem 2.2 with $a = 1$. According to the definition (D.3), the mode d_M is given by

$$d_M = \mu_M + r_M + \sum_{1 \le j < M} H(b_j \otimes b^{(j+1)}_M). \tag{6.13}$$

On the other hand, combining (2.14) and (4.13) with $i = n + 1$, we have $\mathcal{E}_{n+1}(b_1 \otimes \cdots \otimes b_M) = \sum_{1 \le j < m \le M} (\min(\mu_j, \mu_m) - H(b_j \otimes b^{(j+1)}_m))$. (Since $b_1 \otimes \cdots \otimes b_M$ is A_{n-1}-highest, the first term in (4.13) vanishes.) We know $\mathcal{E}_{n+1}(b_1 \otimes \cdots \otimes b_M) = \rho_{n+1}(b_1 \otimes \cdots \otimes b_M)$ by Proposition 4.6. Moreover, since Proposition 6.4 for $A^{(1)}_{n-1}$ is assumed, we are allowed to

use Theorem 4.9 to set $\rho_{n+1}(b_1 \otimes \cdots \otimes b_M) = \tau^{(1)}_{M,n+1}$. Consequently, $\tau^{(1)}_{M,n+1}$ is expressed as

$$\tau^{(1)}_{M,n+1} = \sum_{1 \le j < m \le M} (\min(\mu_j, \mu_m) - H(b_j \otimes b_m^{(j+1)})). \tag{6.14}$$

The formula (6.12) is a corollary of (6.13), (6.14) and the condition $\mu_1 \le \cdots \le \mu_N$. □

Given a rigged configuration $((1^L), (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ with $\mu^{(1)} = (\mu_1, \ldots, \mu_N)$ and $r^{(1)} = (r_1, \ldots, r_N)$, we introduce the numbers

$$k_{M,i} = \min(\mu_{[M]}, \mu_{[M]}) - \min(\mu_{[M-1]}, \mu_{[M-1]}) + r_M + \tau^{(1)}_{M-1,i} - \tau^{(1)}_{M,i} \tag{6.15}$$

for $1 \le M \le N$, $1 \le i \le n+1$.

Lemma 6.7. *Suppose that Proposition 6.4 is true for $A^{(1)}_{n-1}$. Let $((1^L), (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ be a rigged configuration for an asymptotic state. Set $\mu^{(1)} = (\mu_1, \ldots, \mu_N)$ with $\mu_1 \le \cdots \le \mu_N$. Then the following relations are valid:*

$$k_{M,n+1} \le k_{M,n} \le \cdots \le k_{M,2} \le k_{M,1} \quad (1 \le M \le N), \tag{6.16}$$

$$k_{M,1} \le k_{M+1,n+1} \quad (1 \le M \le N-1), \tag{6.17}$$

$$k_{M,1} - k_{M,n+1} = \mu_M \quad (1 \le M \le N). \tag{6.18}$$

Proof. By the assumption we may use Lemma 6.6. The scattering data $b_1[d_1] \otimes \cdots \otimes b_N[d_N]$ considered there should be understood as a normal ordered one here because we deal with an asymptotic state and assume $\mu_1 \le \cdots \le \mu_N$. See the remark before Lemma 6.5. From the definition (6.15), $k_{M,i-1} - k_{M,i} = \tau^{(1)}_{M-1,i-1} - \tau^{(1)}_{M,i-1} - \tau^{(1)}_{M-1,i} + \tau^{(1)}_{M,i}$ for $2 \le i \le n+1$. This is equal to x_i in (6.11) hence nonnegative, proving (6.16). Summing this over $2 \le i \le n+1$ we get (6.18). Comparing (6.12) and (6.15), we have $k_{M,n+1} = d_M + |\mu_{[M-1]}|$. Therefore $k_{M+1,n+1} - k_{M,1} = k_{M+1,n+1} - k_{M,n+1} - \mu_M = d_{M+1} - d_M$. Since d_i's are the modes of normal ordered scattering data, this is nonnegative, showing (6.17). □

Now we are ready to determine the precise form of asymptotic states from the associated rigged configurations.

Lemma 6.8. *Suppose that Proposition 6.4 is true for $A^{(1)}_{n-1}$. For an asymptotic state p, let $((1^L), (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ be its rigged configuration and $\mu^{(1)} = (\mu_1, \ldots, \mu_N)$ with $\mu_1 \le \cdots \le \mu_N$. Then $p = p_1 \otimes \cdots \otimes p_L \in B_1^{\otimes L}$ is given by*

$$p_k = \begin{cases} i & k_{M,i} < k \le k_{M,i-1} \quad (2 \le i \le n+1, 1 \le M \le N), \\ 1 & k_{M,1} < k \le k_{M+1,n+1} \quad (0 \le M \le N), \end{cases} \tag{6.19}$$

where $k_{0,1} = 0$, $k_{N+1,n+1} = L$. Namely p has the form:

$$11 \cdots 11 (b_1) 11 \cdots\cdots 11 (b_M) 11 \cdots 11 (b_{M+1}) 11 \cdots\cdots 11 (b_N) 11 \cdots 11, \tag{6.20}$$

where the segment $(b_M) \in B_1^{\otimes \mu_M}$ (soliton) looks as

$$\begin{array}{ccccccccccc} k_{M,n+1} & & k_{M,n} & k_{M,n-1} & \cdots & k_{M,i} & k_{M,i-1} & \cdots & k_{M,2} & & k_{M,1} \\ \downarrow & & \downarrow & \downarrow & & \downarrow & \downarrow & & \downarrow & & \downarrow \\ n{+}1, \cdots, n{+}1, & n, & \cdots, n, & \cdots\cdots & & , i, & \cdots, i, & \cdots\cdots & , 2, & \cdots, & 2 \end{array} \tag{6.21}$$

Note that Lemma 6.7 guarantees that the regions of k appearing in (6.19) is the disjoint union decomposition of $1 \le k \le L$.

Proof. By the assumption we may use Lemmas 6.6 and 6.7. In particular we use the notation x_i and d_M in Lemma 6.6. Lemma 6.5 tells that p indeed has the form (6.20). The segment (b_M) has the left end at $k' = d_M + |\mu_{[M-1]}| + 1$ and is arranged as $\overbrace{n{+}1\cdots n{+}1}^{x_{n+1}}\cdots\overbrace{2\cdots 2}^{x_2}$ with x_i specified by (6.11). From the proof of Lemma 6.7, we find that $k' = k_{M,n+1} + 1$ and $x_i = k_{M,i-1} - k_{M,i}$. Therefore it looks as (6.21). □

6.3. **Evaluation of ρ_i and τ_i on asymptotic states.** First we evaluate the tau function $\tau_{k,i}$ of asymptotic states in terms of $\tau_i^{(1)}$.

Lemma 6.9. *Suppose that Proposition 6.4 is true for $A_{n-1}^{(1)}$. If $((1^L), (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ is a rigged configuration for an asymptotic state with $\mu^{(1)} = (\mu_1, \ldots, \mu_N)$, $r^{(1)} = (r_1, \ldots, r_N)$, $(\mu_1 \le \cdots \le \mu_N)$, the associated tau function is given by*

$$\tau_{k,i} = Mk - \min(\mu_{[M]}, \mu_{[M]}) - |r_{[M]}| + \tau_{M,i}^{(1)} \quad (k_{M,i} < k \le k_{M+1,i}), \tag{6.22}$$

where $0 \le M \le N, 1 \le i \le n+1$ and $k_{0,i} = 0$, $k_{N+1,i} = L$.

Proof. From (2.23) we know

$$\tau_{k,i} = \max_{\nu \subseteq \mu^{(1)}} \{\ell(\nu)k - \min(\nu, \nu) - |s| + \tau_i^{(1)}(\nu)\}. \tag{6.23}$$

Since s is the rigging attached to ν and runs over the subset of $r^{(1)}$ that satisfies the asymptotic condition (6.3), the choice of ν that attains the maximum must be of the form $\nu = \mu_{[M]}$ for some $0 \le M \le N$. (We interpret $\mu_{[0]} = \emptyset$.) In terms of the notation (2.24), we have $\tau_i^{(1)}(\mu_{[M]}) = \tau_{M,i}^{(1)}$. In (6.23), the quantity in { } at $\nu = \mu_{[M-1]}$ and $\nu = \mu_{[M]}$ become equal if and only if

$$Mk - \min(\mu_{[M]}, \mu_{[M]}) - |r_{[M]}| + \tau_{M,i}^{(1)} = (M \to M-1). \tag{6.24}$$

This yields $k = k_{M,i}$ (6.15). Comparing the k-dependence $(M-1)k$ and Mk, we conclude that $\nu = \mu_{[M]}$ gives a larger value than $\nu = \mu_{[M-1]}$ if $k_{M,i} < k$. Moreover we may use Lemma 6.7 by the assumption and therefore know that $\cdots < k_{M,i} < k_{M+1,i} < \cdots$. Thus we conclude that the maximum in (6.23) is attained at $\nu = \mu_{[M]}$ for $k_{M,i} < k \le k_{M+1,i}$, where $\tau_{k,i}$ is equal to the left hand side of (6.24). □

Next we evaluate $\rho_{k,i}$ for asymptotic states.

Lemma 6.10. *Under the same assumption as Lemma 6.9, $\rho_{k,i}$ for the asymptotic state is given by*

$$\rho_{k,i} = \tau_{k,i} \quad (1 \le k \le L,\ 2 \le i \le n+1), \tag{6.25}$$

where the right hand side is specified by (6.22).

Proof. By the assumption we may use Lemma 6.8, which specifies the concrete form of the asymptotic state as in (6.21). To evaluate $\rho_{k,i}(p)$ (4.1), we count only the balls of colors $2, 3, \ldots, i$ in p itself and those of any color $\{2, \ldots, n+1\}$ in the subsequent states $T_\infty^{t \ge 1}(p)$. From Proposition 3.5 and (6.15), the positions $k_{M,i}$ in (6.21) changes as $k_{M,i} \to k_{M,i} + \mu_M$ under the time evolution. Due to $\mu_1 \le \cdots \le \mu_N$, there is no collision among the segments (solitons) (b_M)'s in (6.20) under the time evolution. In view of these facts, the counting for $\rho_{k,i}$ within the region $k_{M,i} < k \le k_{M+1,i}$ is done as

$$\rho_{k,i} = \sum_{M'=1}^{M} (k - k_{M',i}). \tag{6.26}$$

From (6.15) and Lemma 2.3, this coincides with the right hand side of (6.22). □

Example 6.11. The following figure helps to understand the counting (6.26). Consider an asymptotic state in which the M-th soliton is $(b_M) = 44332$. Its time evolution takes the form:



Here we have omitted $\otimes$, letters 1 and the other solitons for simplicity. Then the contribution to $\rho_{k,3}$ from the M-th soliton comes from the balls within the frame, and their number is certainly equal to $k - k_{M,3}$.

Proof of Proposition 6.4. Due to Lemma 6.10 and induction on n, it now suffices to show $n = 1$ case of Proposition 6.4 to complete its proof. It is Lemma 6.6 that we started relying on the $n - 1$ case. But when $n = 1$, all the subsequent assertions are easily derived by only using Lemma 6.5 and the definitions of the scattering data and normal ordering in Appendix D. In particular, all the formulas are valid by setting $\tau^{(1)}_{M,2} = 0$ and $\tau^{(1)}_{M,1} = -|\mu_{[M]}|$ in agreement with the definition under (2.23). Thus (6.11) becomes $b_M = (x_2)$ with $x_2 = \mu_M$, and (6.12) reads $d_M = |\mu_{[M]}| + r_M$. The definition (6.15) reads $k_{M,2} = k_{M,1} - \mu_M = \min(\mu_{[M]}, \mu_{[M]}) - \min(\mu_{[M-1]}, \mu_{[M-1]}) + r_M$. Using the fact that $r_M \leq r_{M+1}$ for normal ordered scattering data, one can directly verify the properties (6.16)–(6.21). By using them Lemma 6.9 is shown for $n = 1$, and (6.22) reads $\tau_{k,2} = \tau_{k,1} + |\mu_{[M]}| = Mk - \min(\mu_{[M]}, \mu_{[M]}) - |r_{[M]}|$. Finally (6.25) can be checked by substituting the above $k_{M,2}$ into (6.26) with $i = 2$. This proves $n = 1$ case of Proposition 6.4, therefore it is established for any n. □

Summary of proofs. We have finished proving Proposition 6.4. From the arguments in Section 6.1, it leads to Proposition 6.1. Combined with Proposition 5.1, Proposition 6.1 proves Theorem 4.9 as explained in Section 4.4. Combined with (4.5), Theorem 4.9 proves Theorem 2.1.

In the course of these proofs, we have identified the three basic quantities by Proposition 4.6 and Theorem 4.9. The tau function τ_i (2.19) which is a piecewise linear function on the rigged configuration, the CTM for the box-ball system ρ_i (4.1) and the energy $\mathcal{E}_i$ (4.12) . We rephrase it as

Theorem 6.12. *For any rigged configuration* $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ *and the corresponding highest path* $p_1 \otimes \cdots \otimes p_L \in \mathcal{P}_+(\mu^{(0)})$*, the equality*

$$\tau_i(p_1 \otimes \cdots \otimes p_k) = \rho_i(p_1 \otimes \cdots \otimes p_k) = \mathcal{E}_i(p_1 \otimes \cdots \otimes p_k) \tag{6.27}$$

is valid for $1 \leq i \leq n+1$ *and* $1 \leq k \leq L$.

Note that the second equality (Proposition 4.6) has been shown even for non-highest states. The generalization of the first equality to them will be done in Theorem 7.4. Before closing the section we include a few immediate consequences.

Corollary 6.13. *For* $k = L$*, Theorem 6.12 becomes*

$$\tau_i(p_1 \otimes \cdots \otimes p_L) = \rho_i(p_1 \otimes \cdots \otimes p_L) = \mathcal{E}_i(p_1 \otimes \cdots \otimes p_L) = -c(\mu, r) - |\mu^{(i)}|, \tag{6.28}$$

where $c(\mu, r)$ *is the value of (2.20) at the "full choice"* $\forall(\nu^{(a)}, s^{(a)}) = (\mu^{(a)}, r^{(a)})$*, and we employ the convention* $|\mu^{(n+1)}| = 0$ *as in (2.19).*

Proof. For $i = n+1$, the equality $\mathcal{E}_{n+1}(p_1 \otimes \cdots \otimes p_L) = -c(\mu, r)$ is a consequence of the known relation between the charge of rigged configurations and the energy of paths [10, 22]. For i general, we find from (4.1) that $\rho_{n+1}(p_1 \otimes \cdots \otimes p_L) - \rho_i(p_1 \otimes \cdots \otimes p_L)$ is the number of balls with colors $i+1, i+2, \ldots, n+1$ in $p_1 \otimes \cdots \otimes p_L$. By the definition of the KKR bijection, it is equal to $|\mu^{(i)}|$. □

Remark 6.14. Corollary 6.13 tells that if $\lambda = \mu^{(0)}$, the max (2.19) is attained at the full choice $\forall(\nu^{(a)}, s^{(a)}) = (\mu^{(a)}, r^{(a)})$. In particular, (2.23) leads to $\tau^{(a)}_{n+1}(\mu^{(a)}) = \min(\mu^{(a)}, \mu^{(a+1)}) - \min(\mu^{(a+1)}, \mu^{(a+1)}) - |r^{(a+1)}| + \tau^{(a+1)}_{n+1}(\mu^{(a+1)})$.

Now we are able to evaluate the conserved quantity E_l (3.2) for highest states in terms of the rigged configurations.

Proposition 6.15. *Let $p \in \mathcal{P}_+(\mu^{(0)})$ be the highest state corresponding to the rigged configuration $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$. Then, its row transfer matrix energy $E_l(p)$ (3.2) is given by $E_l(p) = \sum_j \min(l, \mu^{(1)}_j)$, which is $E^{(1)}_l$ in (2.8).*

Proof. Combining Proposition 4.8 and Theorem 6.12, we have

$$E_l(p) = \mathcal{E}_{n+1}(p) - \mathcal{E}_{n+1}(T_l(p)) = \tau_{n+1}(\mu^{(0)}) - \tau'_{n+1}(\mu^{(0)}).$$

Here, by Proposition 3.5, $\tau'_{n+1}(\mu^{(0)})$ is obtained from $\tau_{n+1}(\mu^{(0)})$ by replacing the rigging $r^{(1)}_i$ with $r^{'(1)}_i = r^{(1)}_i + \min(l, \mu^{(1)}_i)$. This amounts to changing $-|s|$ in (2.23) (with $a = 0, d = n+1$) into $-|s| - \sum_j \min(l, \nu_j)$. On the other hand from Remark 6.14, we know that the max in (2.23) for $\lambda = \mu^{(0)}$ is attained at $\nu = \mu^{(1)}$. Therefore the difference $\tau_{n+1}(\mu^{(0)}) - \tau'_{n+1}(\mu^{(0)})$ is equal to $\sum_j \min(l, \mu^{(1)}_j)$. □

Proposition 6.15 will be extended to non-highest states in Proposition 7.7.

7. N-soliton solutions of the Box-ball system

As an application of Theorem 2.1, we present the solution of the initial value problem and N-soliton solutions of the box-ball system. To cope with arbitrary states not necessarily highest, we first introduce in Section 7.1 an extension of the rigged configurations for such states, which we expect is equivalent to those studied in [23, 40]. We naturally extend the domain of the tau function to them. Generalizations of Theorems 2.1, 4.9 and 6.12 to arbitrary (non-highest) states are presented in Section 7.2. Based on these results, we give the solution of the initial value problem in Section 7.3. In Section 7.4 we derive several formulas for our tau functions in terms of the parameters that specify solitons. Together with (7.13), they yield the N-soliton solution of the box-ball system. Our approach provides the *general* solution, which accommodates arbitrary number and kinds of solitons. A class of special solutions have been constructed earlier in [26].

7.1. $\boldsymbol{\tau_i}$ for non-highest states.

For $\mu^{(0)} = (\mu^{(0)}_1, \ldots, \mu^{(0)}_L) \in (\mathbb{Z}_{\geq 1})^L$, let $p \in B_{\mu^{(0)}_1} \otimes \cdots \otimes B_{\mu^{(0)}_L}$ be an arbitrary element not necessarily highest. Set

$$\tilde{p} = p_{\mathrm{vac}} \otimes p, \tag{7.1}$$

$$p_{\mathrm{vac}} = (12\ldots n)^{\otimes M_n} \otimes \cdots \otimes (12)^{\otimes M_2} \otimes 1^{\otimes M_1}, \tag{7.2}$$

where $(12\ldots n)$ for example means $1 \otimes \cdots \otimes n \in B_1^{\otimes n}$. The rigged configuration for p_{vac} is given by

$$\mathrm{rc}_{\mathrm{vac}} = ((1^{L_0}), ((1^{L_1}), (0^{L_1})), \ldots, ((1^{L_n}), (0^{L_n}))), \tag{7.3}$$

$$L_a = \sum_{b=1}^{n} (b - \min(a,b)) M_b = \sum_{b=a+1}^{n} (b-a) M_b \quad (0 \leq a \leq n). \tag{7.4}$$

Thus $L_n = 0$ and $((1^{L_n}), (0^{L_n}))$ actually means $(\emptyset, \emptyset)$. The vacancy numbers $p_j^{(a)}$ (2.7) for the configuration $((1^{L_0}), (1^{L_1}), \ldots, (1^{L_n}))$ of $\mathrm{rc}_{\mathrm{vac}}$ is calculated as

$$\delta_{a,1} L_0 - \sum_{b=1}^{n} C_{a,b} L_b = M_a \tag{7.5}$$

for any $j \geq 1$. In (7.1), one can always make the state $\tilde{p}$ highest by taking $M_1, \ldots, M_n$ sufficiently large. In fact, the choice

$$M_a > m_{a+1} \quad (1 \leq a \leq n) \tag{7.6}$$

suffices, where m_a denotes the total number of the letter a contained in the tableau representation of p.

Let $(\tilde{\mu}, \tilde{r}) = (\tilde{\mu}^{(0)}, (\tilde{\mu}^{(1)}, \tilde{r}^{(1)}), \ldots, (\tilde{\mu}^{(n)}, \tilde{r}^{(n)}))$ be the rigged configuration for the highest state $\tilde{p}$. By the definition of the KKR bijection, it "contains" $\mathrm{rc}_{\mathrm{vac}}$ (7.3) for p_{vac}. By this we mean that $(\tilde{\mu}, \tilde{r})$ can be depicted as follows ($n = 3$):



Recall that $\mu^{(0)}$ is not limited to a partition, therefore it is not necessarily a Young diagram. Neither $\tilde{\mu}^{(a)}$ has been depicted so. As mentioned after (2.5), any reordering of $\{(\tilde{\mu}_i^{(a)}, \tilde{r}_i^{(a)})\}$ for each a should be understood as the same rigged configuration.

From the above rigged configuration $(\tilde{\mu}^{(0)}, (\tilde{\mu}^{(1)}, \tilde{r}^{(1)}), \ldots, (\tilde{\mu}^{(n)}, \tilde{r}^{(n)}))$, we extract the data $(\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)})$ by

$$\tilde{\mu}^{(a)} = (\mu_i^{(a)})_{i=1}^{l_a} \sqcup (1^{L_a}), \qquad \mu^{(a)} = (\mu_i^{(a)})_{i=1}^{l_a}, \tag{7.7}$$

$$\tilde{r}^{(a)} = (r_i^{(a)} + M_a)_{i=1}^{l_a} \sqcup (0^{L_a}), \quad r^{(a)} = (r_i^{(a)})_{i=1}^{l_a} \tag{7.8}$$

for $1 \leq a \leq n$, where $l_a = \ell(\mu^{(a)})$. The shift M_a in defining $r_i^{(a)}$ by (7.8) has been introduced on account of (7.5) and the algorithm for the KKR bijection, especially Lemma C.3. As the result, $(\mu^{(1)}, r^{(1)})$, …, $(\mu^{(n)}, r^{(n)})$ become independent of $M_1, \ldots, M_n$ as they get large sufficiently. Therefore the data $(\mu, r) = (\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ is determined unambiguously from $p \in B_{\mu_1^{(0)}} \otimes \cdots \otimes B_{\mu_L^{(0)}}$ by the prescription (7.1)–(7.8). We call (μ, r) the unrestricted rigged configuration for p, which we expect is equivalent to the one studied in [23, 40]. For highest states, it coincides with the rigged configuration under the KKR bijection, but in general $(\mu^{(0)}, \mu^{(1)}, \ldots, \mu^{(n)})$ is not necessarily a configuration.

The vacancy number $p_j^{(a)}$ (2.7) can become negative. The rigging $r^{(a)} \in \mathbb{Z}^{l_a}$ is no longer limited to the range (2.6) but obeys the relaxed condition $r_i^{(a)} \le p^{(a)}_{\mu_i^{(a)}}$ with some non-positive lower bound. We associate the tau function $\tau_d^{(a)}(\lambda)$ ($\lambda \subseteq \mu^{(a)}$) to an unrestricted rigged configuration (μ, r) by the same formula as (2.22). For $\lambda = \mu^{(0)}_{[k]}$ and $p = p_1 \otimes \cdots \otimes p_L$, we will also use the notation $\tau_i(\lambda) = \tau_{k,i} = \tau_i(p_1 \otimes \cdots \otimes p_k)$ as in (2.24).

Example 7.1. Take $n = 3$ and consider the non-highest state p and the highest state $\tilde{p}$ as

$$\begin{aligned} p &= 344 \otimes 2 \otimes 13 \otimes 24 \in B_3 \otimes B_1 \otimes B_2 \otimes B_2, \\ \tilde{p} &= p_{\mathrm{vac}} \otimes p, \\ p_{\mathrm{vac}} &= 123123121 \in B_1^{\otimes 9}, \end{aligned}$$

where we have omitted $\otimes$ in p_{vac}. The rigged configuration $(\tilde{\mu}, \tilde{r})$ for $\tilde{p}$ is



We have

$$(M_1, M_2, M_3) = (1, 1, 2), \quad (L_0, L_1, L_2, L_3) = (9, 5, 2, 0)$$

according to (7.4). Thus the definitions (7.7) and (7.8) yield the unrestricted rigged configuration (μ, r) depicted as



Since $p_3^{(3)} = -2$, this is not a configuration.

7.2. $\boldsymbol{\tau_i = \rho_i}$ for non-highest states.

Lemma 7.2. *For any element $p \in B_{\mu_1^{(0)}} \otimes \cdots \otimes B_{\mu_L^{(0)}}$, let p_{vac}, L_a, $(\tilde{\mu}, \tilde{r})$ and $(\mu, r) = (\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ be as in (7.2)–(7.8). For a fixed $\lambda \subseteq \mu^{(0)}$, the tau function associated with the rigged configuration $(\tilde{\mu}, \tilde{r})$ is decomposed as*

$$\tau_i(\lambda \sqcup (1^{L_0})) = \tau_i(p_{\mathrm{vac}}) + L_1 \ell(\lambda) + \tau_i(\lambda), \tag{7.9}$$

$$\tau_i(p_{\mathrm{vac}}) = L_0 L_1 - \frac{1}{2} \sum_{1 \le a,b \le n} C_{a,b} L_a L_b - L_i \tag{7.10}$$

for sufficiently large $M_1, M_2, \ldots, M_n$. Here $\tau_i(p_{\mathrm{vac}})$ is the tau function for the rigged configuration $\mathrm{rc}_{\mathrm{vac}}$ (7.3). The last term in the right hand side of (7.9) is the tau function (2.19) associated with the unrestricted rigged configuration (μ, r).

Proof. Let us write down the left hand side of (7.9) according to (2.19) and (2.20) as (7.11)

$$\tau_i(\lambda \sqcup (1^{L_0})) = \max_{\tilde{\nu} \subseteq \tilde{\mu}} \Big\{ \min(\lambda \sqcup (1^{L_0}), \tilde{\nu}^{(1)}) - \frac{1}{2} \sum_{a,b} C_{a,b} \min(\tilde{\nu}^{(a)}, \tilde{\nu}^{(b)}) - \sum_a |\tilde{s}^{(a)}| - |\tilde{\nu}^{(i)}| \Big\}.$$

For $M_1, M_2, \ldots, M_n$ sufficiently large, one has $L_0 \gg L_1 \gg \cdots \gg L_{n-1} \gg 1$. In such a circumstance, one can show that the max can be limited to those $\tilde{\nu}^{(a)} \subseteq \tilde{\mu}^{(a)}$ that contain (1^{L_a}) part entirely. Accordingly, we set

$$\tilde{\nu}^{(a)} = \nu^{(a)} \sqcup (1^{L_a}), \qquad \nu^{(a)} \subseteq \mu^{(a)},$$
$$|\tilde{s}^{(a)}| = |s^{(a)}| + M_a \ell(\nu^{(a)}), \quad s^{(a)} \subseteq r^{(a)},$$

taking (7.7) and (7.8) into account. Substituting these forms into (7.11) and using the formula (7.5) and $\min(\nu^{(a)} \sqcup (1^{L_a}), \nu^{(b)} \sqcup (1^{L_b})) = L_a L_b + L_a \ell(\nu^{(b)}) + L_b \ell(\nu^{(a)}) + \min(\nu^{(a)}, \nu^{(b)})$, we obtain (7.9). The expression (7.10) is derived by means of (6.28). □

A decomposition parallel to (7.9) takes place also for ρ_i.

Lemma 7.3. *Under the same setting as Lemma 7.2, set $p = p_1 \otimes \cdots \otimes p_L$ and take $\lambda = \mu^{(0)}_{[k]}$ in the notation (2.2), hence $\ell(\lambda) = k$. Then for $M_1, M_2, \ldots, M_n$ sufficiently large, the following relation is valid:*

$$\rho_i(p_{\mathrm{vac}} \otimes p_1 \otimes \cdots \otimes p_k) = \rho_i(p_{\mathrm{vac}}) + L_1 k + \rho_i(p_1 \otimes \cdots \otimes p_k). \tag{7.12}$$

Proof. In view of $p_{\mathrm{vac}} \in B_1^{\otimes L_0}$, the time evolution of $p_{\mathrm{vac}} \otimes p_1 \otimes \cdots \otimes p_k$ under T_∞ looks as follows ($n = 3$).



On the top row, the length L_0 part is p_{vac} and the length k part is $p_1 \otimes \cdots \otimes p_k$. By the definition (4.1), $\rho_i(p_{\mathrm{vac}} \otimes p_1 \otimes \cdots \otimes p_k)$ is the number of balls with colors $2, \ldots, i$ on the top row and all the balls in the SW quadrant beneath it.

For $M_1, \ldots, M_n$ sufficiently large, one has $L_0 \gg M_1 \gg 1$. Moreover from the time evolution rule in Proposition 3.2, the left segment within p_{vac} with length $L_0 - M_1$ undergoes just a translation to the right by one lattice unit under T_∞. Thus this segment and the hatched region containing the balls are entirely separated by the strip $11 \ldots 11$ of empty boxes with width $M_1 \gg 1$. Therefore $\rho_i(p_{\mathrm{vac}} \otimes p_1 \otimes \cdots \otimes p_k)$ is decomposed into the contributions from p_{vac} (trapezoid in the bottom left), $p_1 \otimes \cdots \otimes p_k$ (hatched region) and the parallelogram in the bottom. By the definition, the first two are equal to $\rho_i(p_{\mathrm{vac}})$ and $\rho_i(p_1 \otimes \cdots \otimes p_k)$, respectively. The last one yields $L_1 k$ because there are L_1 balls in total in the left segment in p_{vac} with length $L_0 - M_1$. □

Now we give the generalization of Theorem 4.9 and Theorem 6.12 to arbitrary (non-highest) states.

Theorem 7.4. *For any state $p = p_1 \otimes \cdots \otimes p_L \in B_{\mu_1^{(0)}} \otimes \cdots \otimes B_{\mu_L^{(0)}}$, let $(\mu, r) = (\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ be the unrestricted rigged configuration, and let τ_i be the associated tau function. Then the equality (6.27), namely, $\tau_i(p_1 \otimes \cdots \otimes p_k) = \rho_i(p_1 \otimes \cdots \otimes p_k) = \mathcal{E}_i(p_1 \otimes \cdots \otimes p_k)$ holds for $1 \le k \le L$.*

Proof. The equality $\rho_i = \mathcal{E}_i$ has been already shown in Proposition 4.6 for any state, and we are only to show $\tau_i = \rho_i$. Since $p_{\mathrm{vac}} \otimes p_1 \otimes \cdots \otimes p_k$ is a highest state associated with the rigged configuration $(\mu^{(0)}_{[k]} \sqcup (1^{L_0}), (\tilde{\mu}^{(1)}, \tilde{r}^{(1)}), \ldots, (\tilde{\mu}^{(n)}, \tilde{r}^{(n)}))$, Theorem 4.9 tells that (7.12) is equal to (7.9) with $\lambda = \mu^{(0)}_{[k]}$. Moreover it also tells that $\tau_i(p_{\mathrm{vac}}) = \rho_i(p_{\mathrm{vac}})$. □

Combining Theorem 7.4 with (4.5), we obtain a generalization of Theorem 2.1 to arbitrary states.

Corollary 7.5. *For any element, $p \in B_{\mu_1^{(0)}} \otimes \cdots \otimes B_{\mu_L^{(0)}}$, let $(\mu, r) = (\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ be the unrestricted rigged configuration. Then $p_k = (x_1, \ldots, x_{n+1}) \in B_{\mu_k^{(0)}}$ is expressed as*

$$x_d = \tau_{k,d} - \tau_{k-1,d} - \tau_{k,d-1} + \tau_{k-1,d-1}$$

in terms of the tau function $\tau_{k,d} = \tau_d((\mu_1^{(0)}, \ldots, \mu_k^{(0)}))$ associated with (μ, r).

7.3. ***N*-soliton solution.** To simplify the notation we write λ in place of $\mu^{(0)}$ in this subsection. We shall exclusively treat the states $p = p_1 \otimes \cdots \otimes p_L \in B_{\lambda_1} \otimes \cdots \otimes B_{\lambda_L}$ such that L is formally infinite and the boundary condition $p_k = u_{\lambda_k}$ is satisfied for $k \gg 1$. Under such a setting, the right hand side of the inequality (7.6) is still finite, therefore all the arguments in Sections 7.1 and 7.2 remain valid.

Our solution of the initial value problem of the box-ball system is formulated as

Theorem 7.6. *For any initial state $p = p_1 \otimes p_2 \otimes \cdots \in B_{\lambda_1} \otimes B_{\lambda_2} \otimes \cdots$, let $(\mu, r) = (\lambda, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ be its unrestricted rigged configuration. Then the state after the time evolution $p'_1 \otimes p'_2 \otimes \cdots = T_{l_1} T_{l_2} \cdots T_{l_t}(p)$ is expressed as $p'_k = (x_1, \ldots, x_{n+1}) \in B_{\lambda_k}$ with*

$$x_d = \tau_{k,d} - \tau_{k-1,d} - \tau_{k,d-1} + \tau_{k-1,d-1}. \tag{7.13}$$

Here $\tau_{k,d} = \tau_d((\lambda_1, \ldots, \lambda_k))$ is the tau function (2.18)–(2.20) associated with $(\lambda, (\mu^{(1)}, r'^{(1)}), (\mu^{(2)}, r^{(2)}), \ldots, (\mu^{(n)}, r^{(n)}))$, where $r_i'^{(1)} = r_i^{(1)} + \sum_{j=1}^{t} \min(l_j, \mu_i^{(1)})$.

Proof. This is a consequence of Corollary 7.5 and Proposition 3.5. □

Let us evaluate the conserved quantity E_l (3.2) in terms of the data $(\lambda, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$.

Proposition 7.7. *For any state $p = p_1 \otimes p_2 \otimes \cdots \in B_{\lambda_1} \otimes B_{\lambda_2} \otimes \cdots$, let $(\mu, r) = (\lambda, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ be its unrestricted rigged configuration. Then the row transfer matrix energy $E_l(p)$ (3.2) is given by $E_l(p) = \sum_j \min(l, \mu_j^{(1)})$.*

When p is highest, this reduces to Proposition 6.15.

Proof. Let $\tilde{p}$ be the highest state (7.1) and let $(\tilde{\mu}, \tilde{r})$ be the corresponding rigged configuration. Proposition 6.15 tells that

$$E_l(\tilde{p}) = \sum_j \min(l, \tilde{\mu}_j^{(1)}) = \sum_j \min(l, (\mu^{(1)} \sqcup (1^{L_1}))_j) = L_1 + \sum_j \min(l, \mu_j^{(1)}),$$

where we have substituted (7.7) into $\tilde{\mu}_j^{(1)}$. On the other hand, due to $M_1 \gg 1$ in (7.2) and the property (3.3), $E_l(\tilde{p})$ is decomposed as $E_l(\tilde{p}) = E_l(p_{\mathrm{vac}}) + E_l(p)$. It is easy to

check $E_l(p_{\mathrm{vac}}) = L_1$ by counting the non-winding number using the graphical rule in Appendix B. □

Following [27, 26, 31], we call those states p of the box-ball system such that $E_l(p) = \sum_{j=1}^{l_1} \min(l, \mu_j^{(1)})$ l_1-soliton states with amplitudes $\mu_1^{(1)}, \dots, \mu_{l_1}^{(1)}$. Thus Proposition 7.7 tells that any state of the box-ball system is an l_1-soliton state for some l_1. Moreover, Theorem 7.6 asserts that in the unrestricted rigged configuration $(\lambda, (\mu^{(1)}, r^{(1)}), \dots, (\mu^{(n)}, r^{(n)}))$, the $A_{n-1}^{(1)}$ part $(\mu^{(1)}, (\mu^{(2)}, r^{(2)}), \dots, (\mu^{(n)}, r^{(n)}))$ is the conserved quantity among which $\mu^{(1)}$ provides the list of amplitudes of solitons. In the remainder of this section we set

$$l_1 = N, \;\; \mu^{(1)} = (\mu_1, \dots, \mu_N), \;\; r^{(1)} = (r_1, \dots, r_N),$$

and rewrite the tau function in terms of the parameters that specify solitons. These parameters are equivalent to the conserved quantity $(\mu, (\mu^{(2)}, r^{(2)}), \dots, (\mu^{(n)}, r^{(n)}))$ as we will see shortly. The result yields the general N-soliton solution of the box-ball system, which supplements the special solution in [26].

From [30, 27, 31], it is known that N-soliton states in the $A_n^{(1)}$ box-ball system are labelled with the $A_{n-1}^{(1)}$ affine crystal $\mathrm{Aff}(B_{\mu_1}^{\geq 2}) \otimes \cdots \otimes \mathrm{Aff}(B_{\mu_N}^{\geq 2})$. The classical part $B_{\mu_1}^{\geq 2} \otimes \cdots \otimes B_{\mu_N}^{\geq 2}$ parametrizes the internal degrees of freedom of solitons. The affine part is incorporated in the integers $r_1, \dots, r_N$, and specifies the positions of the solitons. Thus we start with any such data

$$b_1 \otimes \cdots \otimes b_N \in B_{\mu_1}^{\geq 2} \otimes \cdots \otimes B_{\mu_N}^{\geq 2}, \quad (r_1, \dots, r_N) \in \mathbb{Z}^N, \tag{7.14}$$

where we call each b_i a soliton. Let $(\mu, (\mu^{(2)}, r^{(2)}), \dots, (\mu^{(n)}, r^{(n)}))$ be the unrestricted rigged configuration for $b_1 \otimes \cdots \otimes b_N$. Without loss of generality we assume

$$\mu_1 \leq \cdots \leq \mu_N, \qquad r_i \leq r_j \;\text{ if }\; \mu_i = \mu_j \text{ and } i < j. \tag{7.15}$$

For any $\nu \subseteq \mu$, let us express the $A_{n-1}^{(1)}$ tau function $\tau_i^{(1)}(\nu)$ associated with $(\mu, (\mu^{(2)}, r^{(2)}), \dots, (\mu^{(n)}, r^{(n)}))$ in terms of $b_1, \dots, b_N$. We parametrize $\nu \subseteq (\mu_1, \dots, \mu_N)$ as $\nu = (\mu_{j_1}, \dots, \mu_{j_M})$ in terms of the subset $J = \{j_1 < \cdots < j_M\} \subseteq \{1, 2, \dots, N\}$. From the array of N solitons $b_1 \otimes \cdots \otimes b_N$ we extract an element in $B_{\mu_{j_1}}^{\geq 2} \otimes \cdots \otimes B_{\mu_{j_M}}^{\geq 2}$ by sending the corresponding components to the left by the combinatorial R as follows:

$$\begin{aligned} B_{\mu_1}^{\geq 2} \otimes \cdots\cdots \otimes B_{\mu_N}^{\geq 2} \quad &\simeq \quad B_{\mu_{j_1}}^{\geq 2} \otimes \cdots \otimes B_{\mu_{j_M}}^{\geq 2} \otimes (\cdots) \\ b_1 \otimes \cdots\cdots \otimes b_N \quad &\simeq \quad b_{j_1}^{(1)} \otimes \cdots \otimes b_{j_M}^{(M)} \otimes (\cdots). \end{aligned} \tag{7.16}$$

A caution is necessary about this notation. Consider for instance $N = 3, M = 2$ cases

$$\begin{aligned} b_1 \otimes b_2 \otimes b_3 &\simeq b_1 \otimes b_3^{(2)} \otimes (\cdot) \quad \text{for } J = \{1, 3\}, \\ &\simeq b_2^{(1)} \otimes b_3^{(2)} \otimes (\cdot) \quad \text{for } J = \{2, 3\}. \end{aligned}$$

Obviously, the elements represented by the same symbol $b_3^{(2)}$ in the two lines are not equal in general. In this way, $b_{j_\alpha}^{(\alpha)}$ is uniquely determined only by further specifying J except $\alpha = 1$. In what follows we will always take it for granted that J has been prescribed.

From Theorem 7.4 for $A_{n-1}^{(1)}$, we know

$$\tau_i^{(1)}(\nu) = \mathcal{E}_i(b_{j_1}^{(1)} \otimes \cdots \otimes b_{j_M}^{(M)}) \quad (2 \leq i \leq n+1).$$

Applying the formula (4.13) for $A_{n-1}^{(1)}$ to the right hand side we get

$$\tau_i^{(1)}(\nu) = \sum_{j \in J} (b_{j,3}^{(1)} + b_{j,4}^{(1)} + \cdots + b_{j,i}^{(1)}) + \mathcal{E}_i^{\vee}(b_{j_1}^{(1)} \otimes \cdots \otimes b_{j_M}^{(M)}) \quad (2 \leq i \leq n+1),$$

where $b_j^{(1)} = (b_{j,2}^{(1)}, \dots, b_{j,n+1}^{(1)})$ is the representation in terms of the number of tableau letters as in (2.11). The case $i = 1$ needs an independent derivation. We recall the

definition $\tau_1^{(1)}(\nu) = \tau_{n+1}^{(1)}(\nu) - |\nu|$ given just before (2.22). Substituting the above formula with $i = n+1$ to this, we find the result is unified into the single formula

$$\tau_i^{(1)}(\nu) = |\nu| - \sum_{j\in J}(b_{j,i+1}^{(1)} + \cdots + b_{j,n+1}^{(1)} + b_{j,2}^{(1)}) + \mathcal{E}_i^\vee(b_{j_1}^{(1)} \otimes \cdots \otimes b_{j_M}^{(M)}) \quad (1 \le i \le n+1), \tag{7.17}$$

under the convention

$$\mathcal{E}_1^\vee(b_{j_1}^{(1)} \otimes \cdots \otimes b_{j_M}^{(M)}) = \mathcal{E}_{n+1}^\vee(b_{j_1}^{(1)} \otimes \cdots \otimes b_{j_M}^{(M)}).$$

This is natural in view of the mod n structure of the indices in $A_{n-1}^{(1)}$. Similarly, the sum in (7.17) may well be written as $b_{j,i+1}^{(1)} + \cdots + b_{j,n+2}^{(1)}$.

Now we are ready to express the $A_n^{(1)}$ tau function (2.23) associated with $(\lambda, (\mu, r), (\mu^{(2)}, r^{(2)}), \ldots, (\mu^{(n)}, r^{(n)}))$:

$$\tau_{k,i} = \max_{\nu\subseteq\mu}\{\min(\lambda_{[k]}, \nu) - \min(\nu, \nu) - |s| + \tau_i^{(1)}(\nu)\} \quad (k \ge 1,\, 1 \le i \le n+1) \tag{7.18}$$

in terms of the solitons $b_1 \otimes \cdots \otimes b_N$ and their positions $r_1, \ldots, r_N$. We parametrize ν by $J = \{j_1, \ldots, j_M\} \subseteq \{1, \ldots, N\}$ as before, and introduce the functions:

$$\eta_{k,i}(j) = \min(\lambda_{[k]}, \mu_j) - r_j - (b_{j,i+1}^{(1)} + \cdots + b_{j,n+1}^{(1)} + b_{j,2}^{(1)}) \quad (j \in J), \tag{7.19}$$

$$\Delta_i(J) = 2\sum_{\substack{l,m\in J\\ l<m}} \min(\mu_l, \mu_m) - \mathcal{E}_i^\vee(b_{j_1}^{(1)} \otimes \cdots \otimes b_{j_M}^{(M)}), \tag{7.20}$$

where $\min(\lambda_{[k]}, \mu_j) = \sum_{m=1}^k \min(\lambda_m, \mu_j)$ according to (2.3). (To simplify the formula, $\min(\mu_l, \mu_m)$ has been kept as it is despite (7.15).) Substituting (7.17) into (7.18) and noting that $\min(\nu, \nu) - |\nu| = 2\sum_{l,m\in J, l<m}\min(\mu_l, \mu_m)$ and $|s| = \sum_{j\in J} r_j$, we find that $\tau_{k,i}$ is expressed as

$$\tau_{k,i} = \max_{J\subseteq\{1,\ldots,N\}}\Big\{\sum_{j\in J}\eta_{k,i}(j) - \Delta_i(J)\Big\} \tag{7.21}$$

for $k \ge 1$, $1 \le i \le n+1$. We introduce $\tau_{k,0} = \tau_{k,n+1} - |\lambda_{[k]}|$ according to (2.18). Then by Theorem 7.6, the local states are specified by (7.13) and the time evolution T_l is given by changing r_j to $r_j + \min(l, \mu_j)$, i.e., $\eta_{k,i}(j)$ into $\eta_{k,i}(j) - \min(l, \mu_j)$.

Using the formula (7.21), it is easy to evaluate the local state (7.13) explicitly for $k \gg 1$ if $\lambda_k = 1$ in this region and the condition (6.3) (without the super script "(1)" in the present notation) is satisfied. It yields the asymptotic state of the box-ball system well after the collisions of solitons. Omitting the derivation similar to Lemma 6.8, we give the final result:

$$\cdots 11\cdots 11(b_1)11\cdots\cdots 11(b_M)\overbrace{11\cdots 11}^{w}(b_{M+1})11\cdots\cdots 11(b_N)11\cdots 11\cdots, \tag{7.22}$$

where $1 \in B_1$ and the symbol $\otimes$ has been suppressed. For each $b_M = (x_2, \ldots, x_{n+1}) \in B_{\vec{\mu}_M}^{\ge 2}$, $(b_M) \in B_1^{\otimes\mu_M}$ stands for the array

$$\overbrace{n{+}1\ \ldots\ n{+}1}^{x_{n+1}}\ \overbrace{n\ \ldots\ n}^{x_n}\ \ldots\ldots\ \overbrace{2\ \ldots\ 2}^{x_2}\ .$$

In (7.22), the interval of adjacent solitons is given by $w = r_{M+1} - r_M + \delta$, where δ is a constant independent of $r_1, \ldots, r_N$. Therefore if $\mu_M < \mu_{M+1}$, we have $w \gg 1$ due to $r_M \ll r_{M+1}$. In case $\mu_M = \mu_{M+1}$, we have

$$w = r_{M+1} - r_M + H(b_M \otimes b_{M+1}) \ge H(b_M \otimes b_{M+1}) \tag{7.23}$$

because of (7.15). Here $H(b_M \otimes b_{M+1})$ is the energy (2.14) for $A_{n-1}^{(1)}$ crystals. It is known (cf. [27, 31]) that $H(b_M \otimes b_{M+1})$ is the minimum distance until which the solitons of the

same amplitude can get close. Therefore (7.23) is consistent with the fact that the tau function (7.21) constructed from the data (7.14) covers all the N-soliton solutions.

Our formula (7.21) possesses a structure analogous to the well known tau function of the KP hierarchy [37]. For each J, the sum $\sum_{j\in J}\eta_{k,i}(j)$ is the superposition of individual solitons, whereas the quantity $\Delta_i(J)$ reflects a multi-body effect. A characteristic feature in $\eta_{k,i}(j)$ (7.19) is that it contains $b_j^{(1)}$ in (7.16) rather than b_j that appears in the asymptotic state (7.22). As for $\Delta_i(J)$, using the definition (4.11), it is "factorized" into the two-body function as

$$\Delta_i(J) = \sum_{1\le\beta<\alpha\le M} S_i(b_{j_\beta}^{(\beta)}\otimes b_{j_\alpha}^{(\beta+1)}), \tag{7.24}$$

$$S_i(b\otimes c) = 2\min(l,m) - Q_i(b\otimes c) \quad (b\otimes c\in B_l^{\ge 2}\otimes B_m^{\ge 2}). \tag{7.25}$$

Here $b_{j_\alpha}^{(\beta+1)}$ is determined by sending $b_{j_\alpha}^{(\alpha)}$ in (7.16) to the left by the combinatorial R as

$$\begin{aligned} b_{j_1}^{(1)}\otimes\cdots\otimes b_{j_\beta}^{(\beta)}\otimes\cdots\otimes b_{j_\alpha}^{(\alpha)} &\simeq b_{j_1}^{(1)}\otimes\cdots\otimes b_{j_\beta}^{(\beta)}\otimes\cdots\otimes b_{j_\alpha}^{(\alpha-1)}\otimes(\cdot)\\ &\simeq b_{j_1}^{(1)}\otimes\cdots\otimes b_{j_\beta}^{(\beta)}\otimes b_{j_\alpha}^{(\beta+1)}\otimes(\cdot\cdots\cdot). \end{aligned}$$

S_i in (7.25) is equal to $\min(l,m)$ plus the i th winding number $\min(l,m)-Q_i(b\otimes c)$. For $i=n+1$, it has been identified as the two-body phase shift of the solitons labelled with b and c [27, 31]. Thus $\Delta_i(J)$ can be regarded as a generalization of it to the multi-body phase shift for an arbitrary color i.

7.4. **Alternative forms of N-soliton solution.** We retain the notation in the previous subsection. The N-soliton solution (7.21) has been expressed in terms of the parameters in (7.14). Here we rewrite it further in terms of the scattering data (Appendices D and E):

$$b_1[d_1]\otimes\cdots\otimes b_N[d_N]\in \mathrm{Aff}(b_{\mu_1}^{\ge 2})\otimes\cdots\otimes\mathrm{Aff}(b_{\mu_N}^{\ge 2}), \tag{7.26}$$

$$d_j = r_j + \sum_{0\le k<j} H(b_k\otimes b_j^{(k+1)}), \quad b_0 = 2^l \ (l\gg 1). \tag{7.27}$$

Our task is essentially to switch from the position (rigging) r_j to the mode d_j. See (2.16) for the symbol 2^l. The mode d_j here is a natural generalization of the one defined by (D.3). In fact, when $b_1\otimes\cdots\otimes b_N$ is a highest element with respect to A_{n-1}, one has $b_j^{(1)}=2^{\mu_j}$ and $H(b_0\otimes b_1^{(1)})=\mu_j$, hence (7.27) reduces to (D.3). The mode is transformed according to (A.1) under the combinatorial R. The affinization of (7.16) reads

$$\begin{aligned} \mathrm{Aff}(B_{\mu_1}^{\ge 2})\otimes\cdots\cdots\otimes\mathrm{Aff}(B_{\mu_N}^{\ge 2}) &\simeq \mathrm{Aff}(B_{\mu_{j_1}}^{\ge 2})\otimes\cdots\otimes\mathrm{Aff}(B_{\mu_{j_M}}^{\ge 2})\otimes(\cdots)\\ b_1[d_1]\otimes\cdots\cdots\otimes b_N[d_N] &\simeq b_{j_1}^{(1)}[d_{j_1}^{(1)}]\otimes\cdots\otimes b_{j_M}^{(M)}[d_{j_M}^{(M)}]\otimes(\cdots). \end{aligned} \tag{7.28}$$

For the notation $d_{j_\alpha}^{(\alpha)}$, the same caution as for $b_{j_\alpha}^{(\alpha)}$ is necessary as mentioned under (7.16). Applying the definition (7.27) to $b_{j_1}^{(1)}[d_{j_1}^{(1)}]\otimes\cdots\otimes b_{j_M}^{(M)}[d_{j_M}^{(M)}]$ in the above, we find

$$d_{j_\alpha}^{(\alpha)} = r_{j_\alpha} + \sum_{0\le\beta<\alpha} H(b_{j_\beta}^{(\beta)}\otimes b_{j_\alpha}^{(\beta+1)}),$$

where the notation is the same as (7.24) and we have employed the convention $j_0=0$ and $b_0^{(0)}=b_0$. The element $b_0\in B_l^{\ge 2}$ in (7.27) is the $A_{n-1}^{(1)}$ analogue of u_∞ appearing in (4.12) for $A_n^{(1)}$. By using (4.11), (4.12) and (2.14) for $A_{n-1}^{(1)}$ crystals, this can be rewritten as

$$d_{j_\alpha}^{(\alpha)} = r_{j_\alpha} - \mathcal{E}_{n+1}(b_{j_1}^{(1)}\otimes\cdots\otimes b_{j_\alpha}^{(\alpha)}) + \mathcal{E}_{n+1}(b_{j_1}^{(1)}\otimes\cdots\otimes b_{j_{\alpha-1}}^{(\alpha-1)}) + \sum_{0\le\beta\le\alpha}\min(\mu_{j_\alpha},\mu_{j_\beta}),$$

where $\min(\mu_0, \mu_{j_\beta}) = \mu_{j_\beta}$. Taking the sum over α and using (4.13), we get

$$\sum_{\alpha=1}^{M} d_{j_\alpha}^{(\alpha)} = \sum_{j\in J} r_j + \sum_{j\in J} b_{j,2}^{(1)} - \mathcal{E}_{n+1}^{\vee}(b_{j_1}^{(1)} \otimes \cdots \otimes b_{j_M}^{(M)}) + \sum_{1\le\beta\le\alpha\le M} \min(\mu_{j_\alpha}, \mu_{j_\beta}), \tag{7.29}$$

where we have used $b_{j,2}^{(1)} + \cdots + b_{j,n+1}^{(1)} = \mu_j$. On the other hand, from Corollary 7.5 we deduce

$$\sum_{\alpha=1}^{M} (b_{j_\alpha,2}^{(\alpha)} + \cdots + b_{j_\alpha,i}^{(\alpha)}) = \tau_i^{(1)}(\nu) - \tau_1^{(1)}(\nu) = \tau_i^{(1)}(\nu) - \tau_{n+1}^{(1)}(\nu) + |\nu| \quad (1 \le i \le n+1),$$

where $\nu = (\mu_{j_1}, \ldots, \mu_{j_M})$ as in the previous subsection. Since $\tau_i^{(1)} = \mathcal{E}_i$ for $A_{n-1}^{(1)}$ by Theorem 7.4, the right hand side here is evaluated by using (4.13), leading to

(7.30)

$$\begin{aligned}
&\sum_{\alpha=1}^{M} (b_{j_\alpha,2}^{(\alpha)} + \cdots + b_{j_\alpha,i}^{(\alpha)}) \\
&= -\sum_{j\in J} (b_{j,i+1}^{(1)} + \cdots + b_{j,n+1}^{(1)}) + \mathcal{E}_i^{\vee}(b_{j_1}^{(1)} \otimes \cdots \otimes b_{j_M}^{(M)}) - \mathcal{E}_{n+1}^{\vee}(b_{j_1}^{(1)} \otimes \cdots \otimes b_{j_M}^{(M)}) + |\nu|.
\end{aligned}$$

From (7.29) and (7.30), the quantity appearing in (7.21) is rewritten as

$$\begin{aligned}
&\sum_{j\in J} \eta_{k,i}(j) - \Delta_i(J) = \min(\lambda_{[k]}, \nu) + \sum_{\alpha=1}^{M} (-\phi_{j_\alpha}^{(\alpha)} + b_{j_\alpha,2}^{(\alpha)} + \cdots + b_{j_\alpha,i}^{(\alpha)}), \\
&\phi_{j_\alpha}^{(\alpha)} = d_{j_\alpha}^{(\alpha)} + \mu_{j_\alpha} + \sum_{1\le\beta<\alpha} \min(\mu_{j_\alpha}, \mu_{j_\beta}) = r_{j_\alpha} + \sum_{0\le\beta<\alpha} S_{n+1}(b_{j_\beta} \otimes b_{j_\alpha}^{(\beta+1)}),
\end{aligned}$$

where S_{n+1} is defined in (7.25). Thus we obtain

$$\tau_{k,i} = \max_{J\subseteq\{1,\ldots,N\}} \Big\{ \sum_{\alpha=1}^{M} \big(\min(\lambda_{[k]}, \mu_{j_\alpha}) - \phi_{j_\alpha}^{(\alpha)} + b_{j_\alpha,2}^{(\alpha)} + \cdots + b_{j_\alpha,i}^{(\alpha)} \big) \Big\} \quad (1 \le i \le n+1), \tag{7.31}$$

where the max extends over all the subsets $J = \{j_1, \ldots, j_M\} \subseteq \{1, \ldots, N\}$. Compared with (7.21), the expression (7.31) is formally free from the multi-body effect. It has been absorbed into the quantity $\phi_{j_\alpha}^{(\alpha)}$, which is a shifted mode.

The formula (7.31) is most naturally presented in terms of the "principal picture" of affine crystals rather than the conventional "homogeneous" one. To explain it, let us make a short digression on the principal picture in this paragraph. Recall that an element in the affine $A_n^{(1)}$ crystal $\mathrm{Aff}(B_l)$ is parametrized as $(x_1, \ldots, x_{n+1})[d]$, where $d \in \mathbb{Z}$ and $x_i \in \mathbb{Z}_{\ge 0}$ are to satisfy $x_1 + \cdots + x_{n+1} = l$. See (2.15). We naturally extend x_i to $i \in \mathbb{Z}$ by $x_{i+n+1} = x_i$. Instead of $(x_1, \ldots, x_{n+1})[d]$, the element is also parametrized as $x_i = \theta_{i-1} - \theta_i$ and $d = \theta_0$ in terms of an infinite sequence $\theta = (\theta_i)_{i\in\mathbb{Z}}$ such that

$$\theta_i \in \mathbb{Z}, \quad \theta_{i-1} \ge \theta_i, \quad \theta_i = \theta_{i+n+1} + l \quad \text{for all } i \in \mathbb{Z}. \tag{7.32}$$

The correspondence between $(x_1, \ldots, x_{n+1})[d]$ and θ is bijective. In fact, $\theta_i = d - x_1 - x_2 - \cdots - x_i$ for $i \ge 0$ and $\theta_i = d + x_0 + x_{-1} + \cdots + x_{i+1}$ for $i < 0$. We set $\mathrm{Aff}_{\mathrm{p}}(B_l) = \{\theta = (\theta_i)_{i\in\mathbb{Z}} \mid (7.32)\}$ and call the crystal structure induced on it the *principal picture*. Explicitly, it is given as follows:

$$\tilde{e}_j(\theta) = (\theta_i - \delta_{i,j}^{(n+1)}), \quad \tilde{f}_j(\theta) = (\theta_i + \delta_{i,j}^{(n+1)}) \ \text{ for } \ \theta = (\theta_i),$$

where $\delta_{i,j}^{(n+1)} = 1$ if $i \equiv j \mod n+1$ and 0 otherwise. If the right hand sides break the condition $\theta_{i-1} \ge \theta_i$ in (7.32), they are to be understood as 0. The combinatorial R is

especially simple in the principal picture:

$$\begin{array}{rcl} R\ :\ \mathrm{Aff}_{\mathrm{p}}(B_l)\otimes \mathrm{Aff}_{\mathrm{p}}(B_m) & \longrightarrow & \mathrm{Aff}_{\mathrm{p}}(B_m)\otimes \mathrm{Aff}_{\mathrm{p}}(B_l) \\ (\theta_i)\otimes(\theta'_i) & \longmapsto & (\theta'_i - S_i)\otimes(\theta_i+S_i). \end{array} \tag{7.33}$$

Here $S_i = S_{i+n+1} = S_i(\theta\otimes\theta')$ is defined to be the color i two-body phase shift $S_i(b\otimes c)$ (7.25) for $A_n^{(1)}$ with $b\otimes c\in B_l\otimes B_m$, where b and c are specified by $b=(\theta_{i-1}-\theta_i)_{i=1}^{n+1}$ and $c=(\theta'_{i-1}-\theta'_i)_{i=1}^{n+1}$. From (2.13), S_i reads explicitly as

$$S_i(\theta\otimes\theta') = 2\min(l,m) - \theta_i + \theta'_{i+n+1} - \min_{1\le k\le n+1}\{\theta'_{i+k}-\theta_{i+k-1}\} \tag{7.34}$$

for $\theta\otimes\theta'\in \mathrm{Aff}_{\mathrm{p}}(B_l)\otimes\mathrm{Aff}_{\mathrm{p}}(B_m)$. Observe the compatibility between (7.33) and (2.12). Actually for $i=0$, the rule (7.33) on θ_0,θ'_0 disagrees with the changes of d,d' in (A.1) under the above mentioned identification $\theta_0=d,\theta'_0=d'$, which renders, however, no problem being merely the discrepancy in the normalizations of the energy function. By $\mathrm{Aff}_{\mathrm{p}}$ we mean the crystal structure including the convention specified in (7.33). θ is a generalized phase variable of solitons.

Back to our N-soliton solution, we restart with the principal picture of the scattering data (7.26):

$$\theta_1\otimes\cdots\otimes\theta_N\in \mathrm{Aff}_{\mathrm{p}}(b^{\ge 2}_{\mu_1})\otimes\cdots\otimes\mathrm{Aff}_{\mathrm{p}}(b^{\ge 2}_{\mu_N}). \tag{7.35}$$

Accordingly, (7.28) reads

$$\begin{array}{ccc} \mathrm{Aff}_{\mathrm{p}}(B^{\ge 2}_{\mu_1})\otimes\cdots\otimes\mathrm{Aff}_{\mathrm{p}}(B^{\ge 2}_{\mu_N}) & \simeq & \mathrm{Aff}_{\mathrm{p}}(B^{\ge 2}_{\mu_{j_1}})\otimes\cdots\otimes\mathrm{Aff}_{\mathrm{p}}(B^{\ge 2}_{\mu_{j_M}})\otimes(\cdots) \\ \theta_1\otimes\cdots\otimes\theta_N & \simeq & \theta^{(1)}_{j_1}\otimes\cdots\otimes\theta^{(M)}_{j_M}\otimes(\cdots), \end{array} \tag{7.36}$$

where, again, the notation $\theta^{(\alpha)}_{j_\alpha}$ is unambiguous only combined with $J=\{j_1,\dots,j_M\}$ as cautioned after (7.16). We set $\theta^{(\alpha)}_{j_\alpha}=(\theta^{(\alpha)}_{j_\alpha,i})_{i\in\mathbb{Z}}$ and identify $\theta^{(\alpha)}_{j_\alpha,1}$ with $\phi^{(\alpha)}_{j_\alpha}$ in (7.31). $\theta^{(\alpha)}_{j_\alpha}\in\mathrm{Aff}_{\mathrm{p}}(B^{\ge 2}_{\mu_{j_\alpha}})$ corresponds to $(b^{(\alpha)}_{j_\alpha,2},\dots,b^{(\alpha)}_{j_\alpha,n+1})[\phi^{(\alpha)}_{j_\alpha}]\in\mathrm{Aff}(B^{\ge 2}_{\mu_{j_\alpha}})$. Therefore we have $\theta^{(\alpha)}_{j_\alpha,i}=\phi^{(\alpha)}_{j_\alpha}-b^{(\alpha)}_{j_\alpha,2}-\cdots-b^{(\alpha)}_{j_\alpha,i}$ for $1\le i\le n+1$. In this way (7.31) is simplified to

$$\tau_{k,i} = \max_{J\subseteq\{1,\dots,N\}}\Big\{\sum_{\alpha=1}^{M}\big(\min(\lambda_{[k]},\mu_{j_\alpha})-\theta^{(\alpha)}_{j_\alpha,i}\big)\Big\}\quad (1\le i\le n+1), \tag{7.37}$$

where the max extends over all the subsets $J=\{j_1,\dots,j_M\}\subseteq\{1,\dots,N\}$ as in (7.31). Note that $\theta^{(\alpha)}_{j_\alpha,1}=\theta^{(\alpha)}_{j_\alpha,n+1}+\mu_{j_\alpha}$ is consistent with the time evolution rule in Proposition 3.5 and $\tau_{k,1}(p)=\tau_{k,n+1}(T_\infty(p))$ indicated by (4.2).

Finally we present an operator formalism that formally leads to (7.37) via the ultra-discretization. Let q be an indeterminate. Let $\mathcal{A}$ be the algebra over $\mathbb{C}[q,q^{-1}]$ generated by the symbols $\Psi(\theta),\Psi^*(\theta)$ ($\theta\in\mathrm{Aff}_{\mathrm{p}}(B_l^{\ge 2})$) that satisfy the commutation relations ($\theta\in\mathrm{Aff}_{\mathrm{p}}(B_l^{\ge 2})$, $\theta'\in\mathrm{Aff}_{\mathrm{p}}(B_m^{\ge 2})$):

$$\Psi(\theta)\Psi^*(\theta') = \Psi^*(\tilde\theta')\Psi(\tilde\theta). \tag{7.38}$$

Here $\tilde\theta,\tilde\theta'$ are related to θ,θ' by the combinatorial R (7.33), (7.34):

$$\theta\otimes\theta'\simeq\tilde\theta'\otimes\tilde\theta. \tag{7.39}$$

(The commutation relation of $\Psi(\theta)\Psi(\theta')$ and $\Psi^*(\theta)\Psi^*(\theta')$ are not needed in the sequel.) We equip $\mathcal{A}$ with the time evolution T_l ($l\in\mathbb{Z}_{\ge 1}$):

$$\begin{array}{l} T_l\Psi(\theta)T_l^{-1}=\Psi(T_l(\theta)),\quad T_l\Psi^*(\theta)T_l^{-1}=\Psi^*(T_l(\theta)), \\ T_l(\theta)=(\theta_i+\min(l,m))\ \text{ for }\ \theta=(\theta_i)\in\mathrm{Aff}_{\mathrm{p}}(B_m^{\ge 2}). \end{array} \tag{7.40}$$

T_l is an automorphism of $\mathcal{A}$ since it commutes with the combinatorial R, i.e., $T_l(\theta)\otimes T_l(\theta')\simeq T_l(\tilde\theta')\otimes T_l(\tilde\theta)$ holds under (7.39). Obviously, $T_lT_m=T_mT_l$ is valid.

For $i \in \mathbb{Z}$, let the bracket $\langle\cdot\rangle_i : \mathcal{A} \to \mathbb{C}[q, q^{-1}]$ be the linear form on $\mathcal{A}$ characterized by the following properties:

$$\langle 1\rangle_i = 1, \quad \langle X\Psi(\theta)\rangle_i = \langle X\rangle_i, \quad \langle \Psi^*(\theta)X\rangle_i = q^{\theta_i}\langle X\rangle_i \quad \text{for } \theta = (\theta_i)_{i\in\mathbb{Z}}, \tag{7.41}$$

where X denotes an arbitrary element in $\mathcal{A}$. We shall write $\langle T_l^k X T_l^{-k}\rangle_i$ simply as $\langle T_l^k X\rangle_i$ for any $k \in \mathbb{Z}$. As an example, let $\theta \otimes \phi \otimes \chi \in \mathrm{Aff_p}(B_a^{\geq 2}) \otimes \mathrm{Aff_p}(B_b^{\geq 2}) \otimes \mathrm{Aff_p}(B_c^{\geq 2})$. Then one has

$$\begin{aligned}
&\big\langle T_l\big(\Psi(\theta)+\Psi^*(\theta)\big)\big(\Psi(\phi)+\Psi^*(\phi)\big)\big(\Psi(\chi)+\Psi^*(\chi)\big)\big\rangle_i \\
&= \big\langle T_l\Psi(\theta)\Psi(\phi)\Psi(\chi)\big\rangle_i + \big\langle T_l\Psi^*(\theta)\Psi(\phi)\Psi(\chi)\big\rangle_i + \big\langle T_l\Psi(\theta)\Psi^*(\phi)\Psi(\chi)\big\rangle_i + \big\langle T_l\Psi(\theta)\Psi(\phi)\Psi^*(\chi)\big\rangle_i \\
&+ \big\langle T_l\Psi^*(\theta)\Psi^*(\phi)\Psi(\chi)\big\rangle_i + \big\langle T_l\Psi^*(\theta)\Psi(\phi)\Psi^*(\chi)\big\rangle_i + \big\langle T_l\Psi(\theta)\Psi^*(\phi)\Psi^*(\chi)\big\rangle_i \\
&+ \big\langle T_l\Psi^*(\theta)\Psi^*(\phi)\Psi^*(\chi)\big\rangle_i.
\end{aligned}$$

We need the following reordering of $\theta \otimes \phi \otimes \chi$ by the combinatorial R:

$$\theta \otimes \phi \otimes \chi \simeq \phi^{(1)} \otimes \theta' \otimes \chi \simeq \phi^{(1)} \otimes \chi^{(2)} \otimes \theta'' \simeq \theta \otimes \overline{\chi}^{(2)} \otimes \phi' \simeq \chi^{(1)} \otimes \theta''' \otimes \phi'.$$

See (7.36). As cautioned after (7.16), there are two elements $\chi^{(2)}$ and $\overline{\chi}^{(2)}$ that are relevant to χ under the choices $J = \{2, 3\}$ and $\{1, 3\}$, respectively. In terms of these elements, the above bracket is evaluated as

$$\begin{aligned}
&1 + q^{\min(l,a)+\theta_i} + q^{\min(l,b)+\phi_i^{(1)}} + q^{\min(l,c)+\chi_i^{(1)}} \\
&+ q^{\min(l,a)+\min(l,b)+\theta_i+\phi_i} + q^{\min(l,a)+\min(l,c)+\theta_i+\overline{\chi}_i^{(2)}} + q^{\min(l,b)+\min(l,c)+\phi_i^{(1)}+\chi_i^{(2)}} \\
&+ q^{\min(l,a)+\min(l,b)+\min(l,c)+\theta_i+\phi_i+\chi_i}.
\end{aligned}$$

From the commutation relation (7.38), the characterization of the bracket (7.41) and the definition (7.36), it follows that the tau function (7.37) associated to the scattering data $\theta_1 \otimes \cdots \otimes \theta_N$ (7.35) comes out as the ultradiscretization:

$$\tau_{k,i} = \lim_{\epsilon\to+0} \epsilon \log \big\langle \prod_{j=1}^{k} T_{\lambda_j}^{-1}\big(\Psi(\theta_1)+\Psi^*(\theta_1)\big)\cdots\big(\Psi(\theta_N)+\Psi^*(\theta_N)\big)\big\rangle_i \quad (1 \leq i \leq n+1), \tag{7.42}$$

where ϵ is related to q by $q = e^{-1/\epsilon}$. The bracket is expanded into 2^N terms as in the above example ($N = 3$). In each of them, the list of the positions of Ψ^* specifies the subset $J = \{j_1, \ldots, j_M\} \subseteq \{1, \ldots, N\}$ for the relevant contribution in (7.37). The time evolution of the tau function $\tau_{k,i}(T_l(p))$ is obtained from (7.42) by further inserting the product $\prod_{j=1}^{k} T_{\lambda_j}^{-1}$ of the automorphism (7.40).

Unlike the tau function (5.1) for the KP hierarchy, $\mathcal{A}$ is not the Clifford algebra and it is not known to us whether the Laurent polynomial

$$\big\langle \prod_{j=1}^{k} T_{\lambda_j}^{-1}\big(\Psi(\theta_1)+\Psi^*(\theta_1)\big)\cdots\big(\Psi(\theta_N)+\Psi^*(\theta_N)\big)\big\rangle_i$$

satisfies any sort of bilinear relations. However, the formula (7.42) is a most intrinsic way to present our ultradiscrete tau function. It synthesizes the principal features in the theories of solitons and crystal basis, i.e., the free-fermion like structure and the combinatorial R.

8. Summary

In this paper we have introduced the ultradiscrete tau function and exploited several properties related to the KKR bijection and the box-ball systems.

In Section 2, τ_i is introduced in (2.18)–(2.20) as a piecewise linear function on rigged configurations. The piecewise linear formula for the KKR bijection is stated in Theorem 2.1. After a brief exposition on the box-ball system in Section 3, we have furthermore

introduced ρ_i and $\mathcal{E}_i$ in Section 4. ρ_i in (4.1) is the number of balls in the SW quadrant in the time evolution pattern of the box-ball system. $\mathcal{E}_i$ defined by (4.12) and (4.11) is a sum of local energy function in the affine crystal. The fact $\rho_i = \mathcal{E}_i$ has been shown in Proposition 4.6. The two quantities provide analogues of the corner transfer matrix [1] in complementary viewpoints; ρ_i from the box-ball system and $\mathcal{E}_i$ from the crystal base theory. Theorem 2.1 is a consequence of the further identification $\tau_i = \rho_i = \mathcal{E}_i$ in Theorem 6.12. Sections 5 and 6 are devoted to a proof of this fact. In Section 5, τ_i is shown to emerge as an ultradiscretization of the tau functions of the KP hierarchy (Lemma 5.3) and satisfy the Hirota type bilinear equation (Proposition 5.1). In Section 6, $\tau_i = \rho_i$ is proved on the asymptotic states by induction on the rank (Proposition 6.1 and its reduction in Proposition 6.4). These properties are enough to establish the claim $\tau_i = \rho_i$ everywhere. Section 7 gives the generalization of Theorem 2.1 and Theorem 6.12 to arbitrary (non-highest) states. As an application, the solution of the initial value problem in the box-ball system is given in Theorem 7.6. We have also included the formulas (7.21), (7.37) and (7.42) for general N-soliton solutions. Curiously, they are most elegantly presented in terms of affine crystals in the principal picture introduced in Section 7.4.

Acknowledgments. The authors thank Masato Okado, Anne Schilling, Mark Shimozono and Taichiro Takagi for useful discussion. Y.Y. is supported by Grants-in-Aid for Scientific No.17340047. R.S. is grateful to Miki Wadati for warm encouragement during the study. He is a research fellow of the Japan Society for the Promotion of Science.

Appendix A. Crystals and combinatorial R

The crystals B_l used in the main text are crystal bases of irreducible finite-dimensional representations of a quantum affine algebra $U'_q(\mathfrak{g})$. Let us recall basic facts on them following [11, 12].

Let P be the weight lattice, $\{\alpha_i\}_{0\le i\le n}$ the simple roots, and $\{\Lambda_i\}_{0\le i\le n}$ the fundamental weights of $\mathfrak{g}$. A crystal B is a finite set with weight decomposition $B = \sqcup_{\lambda\in P} B_\lambda$. The Kashiwara operators $\tilde{e}_i, \tilde{f}_i$ $(i = 0, 1, \cdots, n)$ act on B as $\tilde{e}_i : B_\lambda \longrightarrow B_{\lambda+\alpha_i} \sqcup \{0\}$, $\tilde{f}_i : B_\lambda \longrightarrow B_{\lambda-\alpha_i} \sqcup \{0\}$. In particular, these operators are nilpotent. By definition, we have $\tilde{f}_i b = b'$ if and only if $b = \tilde{e}_i b'$. For any $b \in B$, set $\varepsilon_i(b) = \max\{m \ge 0 \mid \tilde{e}_i^m b \ne 0\}$ and $\varphi_i(b) = \max\{m \ge 0 \mid \tilde{f}_i^m b \ne 0\}$. Then we have the weight $\mathrm{wt}b$ of b by $\mathrm{wt}b = \sum_{i=0}^n (\varphi_i(b) - \varepsilon_i(b))\Lambda_i$.

For two crystals B and B', one can define the tensor product $B \otimes B' = \{b \otimes b' \mid b \in B, b' \in B'\}$. The operators $\tilde{e}_i, \tilde{f}_i$ act on $B \otimes B'$ by

$$\begin{aligned}
\tilde{e}_i(b \otimes b') &= \begin{cases} \tilde{e}_i b \otimes b' & \text{if } \varphi_i(b) \ge \varepsilon_i(b') \\ b \otimes \tilde{e}_i b' & \text{if } \varphi_i(b) < \varepsilon_i(b'), \end{cases} \\
\tilde{f}_i(b \otimes b') &= \begin{cases} \tilde{f}_i b \otimes b' & \text{if } \varphi_i(b) > \varepsilon_i(b') \\ b \otimes \tilde{f}_i b' & \text{if } \varphi_i(b) \le \varepsilon_i(b'). \end{cases}
\end{aligned}$$

Here $0 \otimes b'$ and $b \otimes 0$ should be understood as 0. For crystals we are considering, there exists a unique isomorphism $B \otimes B' \xrightarrow{\sim} B' \otimes B$, *i.e.* a unique map which commutes with the action of Kashiwara operators. In particular, it preserves the weight.

For a crystal B we define its affinization $\mathrm{Aff}(B) = \{b[d] \mid d \in \mathbb{Z}, b \in B\}$ by $\tilde{e}_i(b[d]) = (\tilde{e}_i b)[d - \delta_{i0}]$ and $\tilde{f}_i(b[d]) = (\tilde{f}_i b)[d + \delta_{i0}]$. ($b[d]$ here corresponds to $T^{-d}af(b)$ in [12].) The crystal isomorphism $B \otimes B' \xrightarrow{\sim} B' \otimes B$ is lifted up to a map $\mathrm{Aff}(B) \otimes \mathrm{Aff}(B') \xrightarrow{\sim} \mathrm{Aff}(B') \otimes \mathrm{Aff}(B)$ called the combinatorial R. It has the following form:

$$\begin{aligned}
R \ : \ \mathrm{Aff}(B) \otimes \mathrm{Aff}(B') &\longrightarrow \mathrm{Aff}(B') \otimes \mathrm{Aff}(B) \\
b[d] \otimes b'[d'] &\longmapsto \tilde{b}'[d' - H(b \otimes b')] \otimes \tilde{b}[d + H(b \otimes b')],
\end{aligned} \tag{A.1}$$

where $b \otimes b' \mapsto \tilde{b}' \otimes \tilde{b}$ under the isomorphism $B \otimes B' \xrightarrow{\sim} B' \otimes B$. $H(b \otimes b')$ is called the energy function and determined up to an additive constant by

$$H(\tilde{e}_i(b \otimes b')) = \begin{cases} H(b \otimes b') + 1 & \text{if } i = 0,\ \varphi_0(b) \geq \varepsilon_0(b'),\ \varphi_0(\tilde{b}') \geq \varepsilon_0(\tilde{b}), \\ H(b \otimes b') - 1 & \text{if } i = 0,\ \varphi_0(b) < \varepsilon_0(b'),\ \varphi_0(\tilde{b}') < \varepsilon_0(\tilde{b}), \\ H(b \otimes b') & \text{otherwise.} \end{cases}$$

Proposition A.1 (Yang-Baxter equation)**.** *The following equation holds on Aff(B) ⊗ Aff(B′) ⊗ Aff(B″):*

$$(R \otimes 1)(1 \otimes R)(R \otimes 1) = (1 \otimes R)(R \otimes 1)(1 \otimes R).$$

We often write the map R simply by $\simeq$. The combinatorial R is naturally restricted to $B \otimes B'$.

In the main text we are concerned about the crystal B_l corresponding to the l-fold symmetric tensor representation. We normalize the energy function so that

$$\max\{H(b \otimes c) \mid b \otimes c \in B_l \otimes B_m\} = \min(l, m).$$

Under this convention one has $\min\{H(b \otimes c) \mid b \otimes c \in B_l \otimes B_m\} = 0$. When $l = m$, the combinatorial R becomes the identity map on $B_l \otimes B_l$ but still acts non-trivially as $R(x[d] \otimes y[e]) = x[e - H(x \otimes y)] \otimes y[d + H(x \otimes y)]$.

Appendix B. Graphical rule for combinatorial R

Following [17], we introduce a graphical rule to calculate the combinatorial R for $A_n^{(1)}$ and energy function given by (2.12) and (2.14). Given the two elements

$$x = (x_1, x_2, \cdots, x_{n+1}) \in B_k, \quad y = (y_1, y_2, \cdots, y_{n+1}) \in B_l,$$

we draw the following diagram to represent the tensor product $x \otimes y$.



Combinatorial R and the energy function H for $B_k \otimes B_l$ (with $k \geq l$) are calculated by the following rule.

(1) Pick any dot, say $\bullet_a$, in the right column and connect it with a dot $\bullet'_a$ in the left column by a line. The partner $\bullet'_a$ is chosen from the dots which are in the lowest row among all dots whose positions are higher than that of $\bullet_a$. If there is no such dot, we return to the bottom and the partner $\bullet'_a$ is chosen from the dots in the lowest row among all dots. In the latter case, we call such a pair or line "winding".
(2) Repeat the procedure (1) for the remaining unconnected dots $(l-1)$-times.
(3) Action of the combinatorial R is obtained by moving all unpaired dots in the left column to the right horizontally. We do not touch the paired dots during this move.
(4) The energy function H is given by the number of winding pairs.

It is known that the results for the combinatorial R and the energy functions are not affected by the order of making pairs ([17] Propositions 3.15 & 3.17). For more properties, including that the above definition indeed satisfies the axiom, see [17].

Example B.1. The diagram for $\boxed{1233}\otimes\boxed{124}$ is



By moving the unpaired dot (letter 2) in the left column to the right, we obtain

$$\boxed{1233}\otimes\boxed{124}\simeq\boxed{133}\otimes\boxed{1224}.$$

Since we have one winding pair, the energy function is $H\left(\boxed{1233}\otimes\boxed{124}\right)=1$.

For $i\in\mathbb{Z}_{n+1}$, the number of connecting lines that cross the horizontal level of the border between x_i and x_{i+1} is called the ith winding number. The energy function H is the $n+1$ th winding number. The quantity $\min(l,k)-(i$ th winding number) is called the i th non-winding number. It is known that $Q_i(x\otimes y)$ in (2.13) gives the i th non-winding number. By the definition, the winding numbers for $x\otimes y$ and $\tilde{y}\otimes\tilde{x}$ are the same if $x\otimes y\simeq\tilde{y}\otimes\tilde{x}$ by the combinatorial R.

Appendix C. KKR bijection

In order to define the Kerov-Kirillov-Reshetikhin (KKR) bijection, there are two different ways. One is the original combinatorial algorithm [9, 10] explained here, and the other one is an algebraic version [34, 35] which will be treated in Appendix D. Although the both definitions are known to be equivalent, they work complementarily in some aspects. In fact, we use the both definitions case by case in the main text.

C.1. **Definition.** The KKR bijection provides one to one correspondence between the set of rigged configurations and that of highest paths. For a given $A_n^{(1)}$ rigged configuration

$$\mathrm{RC}=\left((\mu_j^{(0)}),\,(\mu_j^{(1)},r_j^{(1)}),\,\cdots,(\mu_j^{(n)},r_j^{(n)})\right), \tag{C.1}$$

we define the KKR procedure $\mathrm{RC}\longmapsto p\in B_{\mu_N^{(0)}}\otimes\cdots\otimes B_{\mu_2^{(0)}}\otimes B_{\mu_1^{(0)}}$, which gives a highest path p. See Section 2.2 for definitions of rigged configurations, vacancy numbers $E_j^{(a)}$ and riggings. The data $\mu^{(0)}$ is called quantum space.

Definition C.1. For a given RC, the image (or path) p of the KKR bijection is obtained by the following procedure.

Step 1: For each row of the quantum space $\mu^{(0)}$, we assign the numbers from 1 to N arbitrarily, and reorder it as

$$\mu^{(0)}=\{\mu_N^{(0)},\cdots,\mu_2^{(0)},\mu_1^{(0)}\}. \tag{C.2}$$

Take row $\mu_1^{(0)}$.

Step 2: We name each box of the row $\mu_1^{(0)}$ as

$$\mu_1^{(0)} = \boxed{\alpha_{l_1}^{(0)} \mid \cdots\cdots \mid \alpha_2^{(0)} \mid \alpha_1^{(0)}}. \tag{C.3}$$

Corresponding to the row $\mu_1^{(0)}$, let p_1 be the array of l_1 empty boxes:

$$p_1 = \boxed{ \mid \cdots\cdots \mid \mid }. \tag{C.4}$$

Starting from the box $\alpha_1^{(0)}$, we recursively choose $\alpha_1^{(i)} \in \mu^{(i)}$ by the following Rule 1:

> **Rule 1:** Assume we have already chosen $\alpha_1^{(i-1)} \in \mu^{(i-1)}$. Let $g^{(i)}$ be the set of all the rows of $\mu^{(i)}$ whose lengths w satisfy
>
> $$w \geq col(\alpha_1^{(i-1)}),$$
>
> where the right hand side means the number of columns in $\mu^{(i-1)}$ that are not located to the right of the box $\alpha_1^{(i-1)}$.
>
> Let $g_s^{(i)}$ ($\subset g^{(i)}$) be the set of all the *singular rows* ($\overset{\text{def}}{\Longleftrightarrow}$ rows whose corresponding vacancy number and rigging are equal) in the set $g^{(i)}$. If $g_s^{(i)} \neq \emptyset$, then choose one of the shortest rows of $g_s^{(i)}$, and denote its rightmost box by $\alpha_1^{(i)}$. If $g_s^{(i)} = \emptyset$, then we take $\alpha_1^{(i)} = \cdots = \alpha_1^{(n)} = \emptyset$.

Step 3: From RC, remove boxes $\alpha_1^{(0)}$, $\alpha_1^{(1)}$, $\cdots$, $\alpha_1^{(j_1-1)}$ chosen above, where $j_1 - 1$ is the maximum k such that $\alpha_1^{(k)} \neq \emptyset$. After the removal, construct a new RC by

> **Rule 2:** Calculate the vacancy numbers $p_i^{(a)} = E_i^{(a-1)} - 2E_i^{(a)} + E_i^{(a+1)}$ along the configuration after the removal. For those rows shortened by the removal, assign their vacancy numbers equal to the new riggings. For the other row, keep the original rigging before Step 3.

Put letter j_1 into the leftmost empty box of p_1 as

$$p_1 = \boxed{j_1 \mid \mid \cdots\cdots \mid }. \tag{C.5}$$

Step 4: Repeat Step 2 and Step 3 for the rest of the boxes $\alpha_2^{(0)}$, $\alpha_3^{(0)}$, $\cdots$, $\alpha_{l_1}^{(0)}$ in this order. Put letters j_k into empty boxes of p_1 from left to right.
Step 5: Repeat Step 1 to Step 4 for the rest of the rows $\mu_2^{(0)}$, $\mu_3^{(0)}$, $\cdots$, $\mu_N^{(0)}$ in this order. Then we obtain p_k from $\mu_k^{(0)}$, which we identify with the tableau representation of the element in $B_{\mu_k^{(0)}}$. The image of the KKR bijection is given by $p = p_N \otimes \cdots \otimes p_2 \otimes p_1$. ∎

The above procedure gives a map from rigged configurations to highest paths. Its inverse also admits a similar description. See Theorem 2 of [9].

C.2. **Example of the KKR bijection.** Let us illustrate a typical example of the KKR bijection. For a later convenience, we treat the single column type quantum space. The procedure for general quantum space is quite similar.

Example C.2. We show that the following rigged configuration corresponds to a path $p = 11112221322433$.



In the above diagram, we have specified the boxes to be removed by Step 3 with the symbol "×". Note that the boxes with "×" are the rightmost boxes of the shortest possible singular rows, and their column coordinates are increasing from the left to the right. We can remove three boxes at a time, thus resulting part of a path is $\boxed{3}$. Similarly we can proceed as



$$
\begin{array}{lccccc}
(1^7) & 1 & \boxed{\times}\boxed{\times}\boxed{\times} & 1 & \emptyset & \emptyset \\
 & & \Big\downarrow \boxed{2}^{\otimes 3} & & & \\
(1^4) & & \emptyset & & \emptyset & \emptyset \\
 & & \Big\downarrow \boxed{1}^{\otimes 4} & & & \\
\emptyset & & \emptyset & & \emptyset & \emptyset
\end{array}
$$

By removing all the boxes, we end up with

$$p = \boxed{1}\otimes\boxed{1}\otimes\boxed{1}\otimes\boxed{1}\otimes\boxed{2}\otimes\boxed{2}\otimes\boxed{2}\otimes\boxed{1}\otimes\boxed{3}\otimes\boxed{2}\otimes\boxed{2}\otimes\boxed{4}\otimes\boxed{3}\otimes\boxed{3}\,.$$

The following lemma is useful.

Lemma C.3. *Let $p \in \mathcal{P}_+(\mu^{(0)})$ and $q \in \mathcal{P}_+(\nu^{(0)})$ be the highest paths corresponding to the rigged configurations $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ and $(\nu^{(0)}, (\nu^{(1)}, s^{(1)}), \ldots, (\nu^{(n)}, s^{(n)}))$, respectively. Then the rigged configuration for the highest path $p\otimes q$ is given by*

$$(\mu^{(0)} \sqcup \nu^{(0)}, (\mu^{(1)}, r^{(1)}) \sqcup (\nu^{(1)}, s^{'(1)}), \ldots, (\mu^{(n)}, r^{(n)}) \sqcup (\nu^{(n)}, s^{'(n)})). \tag{C.6}$$

Here $(\nu^{(a)}, s^{'(a)}) = \{(\nu_i^{(a)}, s_i^{'(a)})\}$ and the rigging $s^{'(a)} = (s_i^{'(a)})$ is given by

$$s_i^{'(a)} = s_i^{(a)} + p^{(a)}_{\nu_i^{(a)}},$$

where $p_j^{(a)}$ is the vacancy number (2.7) for $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$.

Proof. Let $q_j^{(a)}$ be the vacancy number for $(\nu^{(0)}, (\nu^{(1)}, s^{(1)}), \ldots, (\nu^{(n)}, s^{(n)}))$. Then the vacancy number $p_j^{'(a)}$ for (C.6) reads $p_j^{'(a)} = p_j^{(a)} + q_j^{(a)}$. Therefore the co-rigging (:= vacancy number − rigging) of the row $(\nu_i^{(a)}, s_i^{'(a)})$ in (C.6) is $p_j^{'(a)} - s_i^{'(a)} = q_j^{(a)} - s_i^{(a)}$ with $j = \nu_i^{(a)}$, which is nothing but the co-rigging of the same row in $(\nu^{(0)}, (\nu^{(1)}, s^{(1)}), \ldots, (\nu^{(n)}, s^{(n)}))$. Recall that the KKR procedure (Definition C.1) consults co-riggings to decide boxes to be removed from a rigged configuration. Therefore the above coincidence of the co-rigging means that the KKR procedure on (C.6) gives the path q when the part $\nu^{(0)}$ is firstly removed from $\mu^{(0)} \sqcup \nu^{(0)}$. Moreover at this stage, the remaining rigged configuration is exactly $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$. □

Appendix D. Vertex operator formalism of the KKR bijection

Here we give a crystal theoretic reformulation of the KKR bijection based on [34, 35]. The central notions are scattering data, normal ordering and the vertex operator. For illustrative examples, see Appendix E.

D.1. **Scattering data and normal ordering.** We call elements of affine crystals $b_1[d_1]\otimes\cdots\otimes b_m[d_m] \in \mathrm{Aff}(B_{l_1})\otimes\cdots\otimes\mathrm{Aff}(B_{l_m})$ *scattering data*. The number d_i is called the i-th mode. By using the combinatorial R, scattering data can be reordered and the modes are changed accordingly. Given a scattering data $s \in \mathrm{Aff}(B_{l_1})\otimes\cdots\otimes\mathrm{Aff}(B_{l_m})$, define $\mathcal{S}_m$ to be the set of such reordering as

$$\mathcal{S}_m = \{s' \in \bigsqcup_{\sigma\in\mathfrak{S}_m}{}' \mathrm{Aff}(B_{l_{\sigma(1)}})\otimes\cdots\otimes\mathrm{Aff}(B_{l_{\sigma(m)}}) \mid s' \simeq s\},$$

where $\sqcup'$ means the disjoint union over all the distinct permutations of $(l_1, \ldots, l_m)$. For instance, if $s = \boxed{234}_7 \otimes \boxed{223}_2$, we have

$$\mathcal{S}_2 = \{\boxed{234}_7 \otimes \boxed{223}_2,\ \ \boxed{234}_0 \otimes \boxed{223}_9\}.$$

Note that in this case, the union over σ is trivial as $(l_1, l_2) = (l_2, l_1) = (3, 3)$, but $\mathcal{S}_2$ contains two distinct elements since the combinatorial R is nontrivial as remarked in the end of Appendix A.

For $i = 2, \dots, m$, let $\mathcal{S}_{i-1}$ be the subset of $\mathcal{S}_i$ having the maximal i-th mode. Then we have

$$\emptyset \neq \mathcal{S}_1 \subseteq \mathcal{S}_2 \subseteq \cdots \subseteq \mathcal{S}_m. \tag{D.1}$$

In the above example, we have $\mathcal{S}_1 = \{\boxed{234}_0 \otimes \boxed{223}_9\}$. We call the elements of $\mathcal{S}_1$ *normal ordered forms* of s. In general the normal ordered form $b_1[d_1] \otimes \cdots \otimes b_m[d_m]$ is not unique but the mode sequence $d_1, \dots, d_m$ is unique by the definition and satisfies $d_1 \leq \cdots \leq d_m$. Any element of $\mathcal{S}_1$ is denoted by $:s:$.

D.2. **Maps $\mathcal{C}^{(1)}, \dots, \mathcal{C}^{(n)}$.** Let $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \dots, (\mu^{(n)}, r^{(n)}))$ be an $A_n^{(1)}$ rigged configuration. Pick the color a part $(\mu^{(a)}, r^{(a)})$. Here we simply write it as (μ, r). Namely $\mu = (\mu_1, \dots, \mu_m)$ is an array of positive integers and $r = (r_i)$, where r_i is the rigging attached to the i-th row in μ of length μ_i. For $1 \leq a \leq n$, let $B_l = B_l^{\geq a+1}$ be the $A_{n-a}^{(1)}$ crystal in the sense explained around (2.17). Define the map $\mathcal{C}^{(a)}$ among the $A_{n-a}^{(1)}$ crystals by

$$\begin{aligned} \mathcal{C}^{(a)} : B_{\mu_1} \otimes \cdots \otimes B_{\mu_m} &\to : \mathrm{Aff}(B_{\mu_1}) \otimes \cdots \otimes \mathrm{Aff}(B_{\mu_m}) : \quad (1 \leq a \leq n) \\ b_1 \otimes \cdots \otimes b_m &\mapsto :b_1[d_1] \otimes \cdots \otimes b_m[d_m]: \end{aligned} \tag{D.2}$$

$$d_i = r_i + \mu_i + \sum_{1 \leq k < i} H(b_k \otimes b_i^{(k+1)}). \tag{D.3}$$

Here $b_i^{(j)} \in B_{\mu_i}$ $(j \leq i)$ is defined by bringing b_i to the left by the combinatorial R as

$$(b_j \otimes \cdots \otimes b_{i-1}) \otimes b_i \simeq b_i^{(j)} \otimes (\cdots)$$

under the isomorphism $(B_{\mu_j} \otimes \cdots \otimes B_{\mu_{i-1}}) \otimes B_{\mu_i} \simeq B_{\mu_i} \otimes (B_{\mu_j} \otimes \cdots \otimes B_{\mu_{i-1}})$. Note that the choice (D.3) is compatible with (A.1).

The map $\mathcal{C}^{(n)}$ involves "$A_0^{(1)}$ crystal" $B_l^{\geq n+1} = \{(n+1)^l\}$. See (2.16) for the notation a^l. The following suffices to define $\mathcal{C}^{(n)}$:

$$(n+1)^l \otimes (n+1)^m \simeq (n+1)^m \otimes (n+1)^l, \quad H((n+1)^l \otimes (n+1)^m) = \min(l, m).$$

Since the normal ordering in (D.2) is not unique, $\mathcal{C}^{(a)}$ is actually multi-valued in general. Here we mean by $\mathcal{C}^{(a)}(\cdot)$ to pick any one of the normal ordered forms. $\mathcal{C}^{(a)}$ is an operator that transforms elements of classical $A_{n-a}^{(1)}$ crystals to normal ordered scattering data by assigning the modes.

D.3. **Maps $\Phi^{(1)}, \dots, \Phi^{(n)}$.** Pick the color a and $a-1$ parts of the configuration and denote them simply by $\mu^{(a)} = (\mu_1, \dots, \mu_m)$ and $\mu^{(a-1)} = (\lambda_1, \dots, \lambda_k)$. Set $B_l = B_l^{\geq a+1}$ and $B'_l = B_l^{\geq a}$. We define the map $\Phi^{(a)}$ from the normal ordered scattering data in $A_{n-a}^{(1)}$ affine crystals to classical $A_{n-a+1}^{(1)}$ crystals:

$$\begin{aligned} \Phi^{(a)} : : \mathrm{Aff}(B_{\mu_1}) \otimes \cdots \otimes \mathrm{Aff}(B_{\mu_m}) : &\to B'_{\lambda_1} \otimes \cdots \otimes B'_{\lambda_k} \quad (1 \leq a \leq n) \\ b_1[d_1] \otimes \cdots \otimes b_m[d_m] &\mapsto c_1 \otimes \cdots \otimes c_k. \end{aligned} \tag{D.4}$$

From (D.3) and the fact that $b_1[d_1] \otimes \cdots \otimes b_m[d_m]$ is normal ordered, we have $0 \leq d_1 \leq \cdots \leq d_m$. Then the image $c_1 \otimes \cdots \otimes c_k$ is determined by the following relation under the isomorphism of $A_{n-a+1}^{(1)}$ crystals: (We write $\mathcal{T}_a^d = \boxed{a}^{\otimes d} \in (B_1^{\geq a})^{\otimes d}$ for short.)

$$\begin{aligned} &(\mathcal{T}_a^{d_1} \otimes b_1 \otimes \mathcal{T}_a^{d_2 - d_1} \otimes b_2 \otimes \cdots \otimes \mathcal{T}_a^{d_m - d_{m-1}} \otimes b_m) \otimes (a^{\lambda_1} \otimes a^{\lambda_2} \otimes \cdots \otimes a^{\lambda_k}) \\ &\simeq (c_1 \otimes \cdots \otimes c_k) \otimes \mathrm{tail}, \end{aligned} \tag{D.5}$$

Here we are regarding $b_i \in B_{\mu_i} = B^{\geq a+1}_{\mu_i}$ as an element of $B'_{\mu_i} = B^{\geq a}_{\mu_i}$ by the natural embedding (2.17) as sets. The tail part has the same structure as $(\mathcal{T}_a^{d_1} \otimes b_1 \otimes \mathcal{T}_a^{d_2-d_1} \otimes \cdots \otimes b_m)$ on the left hand side. In the actual use, it turns out to be $(\mathcal{T}_a^{d_1} \otimes a^{\mu_1} \otimes \mathcal{T}_a^{d_2-d_1} \otimes \cdots \otimes a^{\mu_m})$ containing the letter a only. (This fact will not be used.)

To obtain $c_1 \otimes \cdots \otimes c_k$ using (D.5), one applies the $A^{(1)}_{n-a+1}$ combinatorial R many times to carry $(\mathcal{T}_a^{d_1} \otimes b_1 \otimes \cdots \otimes \mathcal{T}_a^{d_m - d_{m-1}} \otimes b_m)$ through $(a^{\lambda_1} \otimes a^{\lambda_2} \otimes \cdots \otimes a^{\lambda_k})$ to the right. The procedure is depicted as



D.4. **Vertex operator formalism.** Define the $A^{(1)}_0$ crystal element

$$p^{(n)} = (n+1)^{\mu_1^{(n)}} \otimes \cdots \otimes (n+1)^{\mu_{l_n}^{(n)}}. \tag{D.6}$$

Theorem D.1. *The image p of the rigged configuration $(\mu^{(0)}, (\mu^{(1)}, r^{(1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ under the KKR bijection is given by*

$$p = \Phi^{(1)}\mathcal{C}^{(1)}\Phi^{(2)}\mathcal{C}^{(2)} \cdots \Phi^{(n)}\mathcal{C}^{(n)}(p^{(n)}). \tag{D.7}$$

This is announced in [34] and proved in [35]. The theorem asserts that the right hand side is independent of the choices of the possibly non-unique normal ordered forms when applying the maps $\mathcal{C}^{(1)}, \ldots, \mathcal{C}^{(n)}$.

Set

$$p^{(a)} = \Phi^{(a+1)}\mathcal{C}^{(a+1)} \cdots \Phi^{(n)}\mathcal{C}^{(n)}(p^{(n)}) \quad (0 \leq a \leq n-1), \tag{D.8}$$

which belongs to the $A^{(1)}_{n-a}$ crystal $B^{\geq a+1}_{\mu_1^{(a)}} \otimes \cdots \otimes B^{\geq a+1}_{\mu_{l_a}^{(a)}}$. Thus p in (D.7) is $p^{(0)}$.

Corollary D.2. *For $0 \leq a \leq n-1$, $p^{(a)}$ coincides with the image of the truncated rigged configuration $(\mu^{(a)}, (\mu^{(a+1)}, r^{(a+1)}), \ldots, (\mu^{(n)}, r^{(n)}))$ under the KKR bijection.*

By the construction, $\mathcal{C}^{(a)}(p^{(a)})$ is a normal ordered scattering data which is an element of an $A^{(1)}_{n-a}$ affine crystal. Then the map $\Phi^{(a)}$ produces an A_{n-a+1} highest path by injecting the scattering data $\mathcal{C}^{(a)}(p^{(a)})$ into the vacuum state $a^{\mu_1^{(a-1)}} \otimes \cdots \otimes a^{\mu_{l_{a-1}}^{(a-1)}}$. We call $\Phi^{(a)}$ *vertex operator* in this sense. The construction (D.8) involves the family of scattering data and vertex operators for crystals of $A^{(1)}_0 \subset A^{(1)}_1 \subset \cdots \subset A^{(1)}_n$. It can be regarded as a crystal theoretical formulation of the nested Bethe ansatz due to Schultz [41].

Appendix E. Inverse scattering formalism of Box-ball system

This appendix is an exposition of the inverse scattering formalism of the box-ball system mentioned in Section 3.2. We illustrate the calculations of scattering data, normal ordering and vertex operators explained in Appendix D along several examples.

E.1. **Time evolution, scattering data and normal ordering.**

Example E.1. Consider the rigged configuration in Example C.2. We put many $1 \in B_1$ on the both sides of the corresponding path $p = 11112221322433$, and consider its time evolution under T_∞ of the box-ball system. See Section 3.1 for the definition of T_∞.

$$
\begin{aligned}
t=0:&\ 11112222111111332111431111111111111111111111111111111\\
t=1:&\ 11111111222211111332114311111111111111111111111111111\\
t=2:&\ 11111111111122222111133214311111111111111111111111111\\
t=3:&\ 11111111111111112222111332431111111111111111111111111\\
t=4:&\ 11111111111111111111222211324331111111111111111111111\\
t=5:&\ 11111111111111111111111222132243311111111111111111111\\
t=6:&\ 11111111111111111111111112211322433211111111111111111\\
t=7:&\ 11111111111111111111111111221113221433211111111111\\
t=8:&\ 11111111111111111111111111112211113221143321111111\\
t=9:&\ 11111111111111111111111111111122111113221114332111
\end{aligned}
$$

Here the length of the paths is 52, and $t = 5$ state contains the original path as $1^{\otimes 20} \otimes p \otimes 1^{\otimes 18}$. The following rigged configurations correspond to the above paths at each time.



The linear dependence of the rigging on t is in agreement with Proposition 3.5. The following is the list of all the normal ordered scattering data corresponding to each time t of the above paths.

$t = 0, 1, 2, 3, 4$	$\boxed{2222}_{4+4t} \otimes \boxed{233}_{11+3t} \otimes \boxed{34}_{16+2t}$
$t = 5$	$\boxed{2222}_{24} \otimes \boxed{233}_{26} \otimes \boxed{34}_{26}$ $\boxed{2222}_{24} \otimes \boxed{23}_{26} \otimes \boxed{334}_{26}$
$t = 6$	$\boxed{22}_{27} \otimes \boxed{2223}_{29} \otimes \boxed{334}_{29}$ $\boxed{22}_{27} \otimes \boxed{223}_{29} \otimes \boxed{2334}_{29}$
$t = 7, 8, 9$	$\boxed{22}_{15+2t} \otimes \boxed{223}_{11+3t} \otimes \boxed{2334}_{5+4t}$

Compare this list with the above time evolution pattern. Each tensor product component of the scattering data corresponds to a soliton in the path. When the modes of the scattering data are well separated, the normal ordering is unique, and the corresponding path consists of well separated solitons that contain the tableau letters in the scattering data (in the reverse order). $t \neq 5, 6$ are such cases. From the viewpoint of the scattering data, collisions of solitons happen when the modes get close and the normal ordering becomes non-unique. $t = 5, 6$ are such cases. See also Example 2.4 for the tau functions at $t = 5$, where $\tau_{k,i}$ there is relevant to $\tau_{k+20,i}$ here.

Let us illustrate the derivation of the normal ordered scattering data at $t = 5$. At $t = 5$, riggings of $\mu^{(1)}$ attached to the rows of length 2, 3 and 4 are $r_1 = 23$, $r_2 = 22$ and $r_3 = 20$, respectively. By Theorem D.1 and (D.8), we know that $p = \Phi^{(1)}\mathcal{C}^{(1)}(p^{(1)})$, where $\mathcal{C}^{(1)}(p^{(1)})$ is the normal ordered scattering data. It is constructed from the A_2-highest path $p^{(1)}$ containing the letters 2, 3 and 4. According to Corollary D.2, $p^{(1)}$ is the image of the KKR bijection of the following part of the original rigged configuration:



Here $\mu^{(1)}$ plays the role of the quantum space, and the KKR bijection is $A_2^{(1)}$ type with letters 2, 3 and 4. For example, if we can remove only a box from $\mu^{(1)}$, then we have the letter 2 as a part of the path, whereas if boxes are removed from $\mu^{(1)}$, $\mu^{(2)}$ and $\mu^{(3)}$, the letter is 4. Removing the rows of $\mu^{(1)}$ from the top, we obtain the A_2 highest path:

$$p^{(1)} = b_1 \otimes b_2 \otimes b_3 = \boxed{22} \otimes \boxed{223} \otimes \boxed{2334}. \tag{E.1}$$

Assigning this with the modes according to (D.2) and (D.3), we get

$$b_1[d_1] \otimes b_2[d_2] \otimes b_3[d_3] = \boxed{22}_{25} \otimes \boxed{223}_{26} \otimes \boxed{2334}_{25}.$$

To derive the mode $d_3 = 25$, for instance, we calculate $\sum_{1 \le k < 3} H(b_k \otimes b_3^{(k+1)})$ in (D.3) as

$$\boxed{22} \otimes \boxed{223} \overset{0}{\otimes} \boxed{2334} \simeq \boxed{22} \overset{1}{\otimes} \boxed{2223} \otimes \boxed{334},$$

where $a \overset{H}{\otimes} b$ signifies the value of the energy function $H(a \otimes b) = H$. Since in (D.3), we have $r_3 = 20$ and $\mu_3 = 4$, the mode is $d_3 = 20 + 4 + 0 + 1 = 25$.

To find the normal ordered scattering data $\mathcal{C}^{(1)}(p^{(1)})$, we follow the procedure (D.1) and list the following sets:

$$\begin{aligned}
\mathcal{S}_3 &= \{\boxed{22}_{25} \otimes \boxed{223}_{26} \otimes \boxed{2334}_{25}, \boxed{22}_{25} \otimes \boxed{2223}_{25} \otimes \boxed{334}_{26}, \\
&\quad \boxed{222}_{25} \otimes \boxed{23}_{26} \otimes \boxed{2334}_{25}, \boxed{222}_{25} \otimes \boxed{2233}_{25} \otimes \boxed{34}_{26}, \\
&\quad \boxed{2222}_{24} \otimes \boxed{23}_{26} \otimes \boxed{334}_{26}, \boxed{2222}_{24} \otimes \boxed{233}_{26} \otimes \boxed{34}_{26}\}, \\
\mathcal{S}_2 &= \{\boxed{22}_{25} \otimes \boxed{2223}_{25} \otimes \boxed{334}_{26}, \boxed{222}_{25} \otimes \boxed{2233}_{25} \otimes \boxed{34}_{26}, \\
&\quad \boxed{2222}_{24} \otimes \boxed{23}_{26} \otimes \boxed{334}_{26}, \boxed{2222}_{24} \otimes \boxed{233}_{26} \otimes \boxed{34}_{26}\}, \\
\mathcal{S}_1 &= \{\boxed{2222}_{24} \otimes \boxed{23}_{26} \otimes \boxed{334}_{26}, \boxed{2222}_{24} \otimes \boxed{233}_{26} \otimes \boxed{34}_{26}\}.
\end{aligned}$$

The both elements in $\mathcal{S}_1$ serve as the normal ordered scattering data in agreement with the previous list at $t = 5$.

Example E.2. Here is a more intriguing example.



The normal ordered scattering data are listed below.

$t=0,1,2,3$	$\boxed{2222}_{4+4t} \otimes \boxed{233}_{9+3t} \otimes \boxed{34}_{13+2t}$
$t=4$	$\boxed{2222}_{20} \otimes \boxed{233}_{21} \otimes \boxed{34}_{21}$
	$\boxed{2222}_{20} \otimes \boxed{23}_{21} \otimes \boxed{334}_{21}$
	$\boxed{222}_{20} \otimes \boxed{2233}_{21} \otimes \boxed{34}_{21}$
	$\boxed{222}_{20} \otimes \boxed{23}_{21} \otimes \boxed{2334}_{21}$
	$\boxed{22}_{20} \otimes \boxed{2223}_{21} \otimes \boxed{334}_{21}$
	$\boxed{22}_{20} \otimes \boxed{223}_{21} \otimes \boxed{2334}_{21}$
$t=5,6,7,8,9$	$\boxed{22}_{12+2t} \otimes \boxed{223}_{9+3t} \otimes \boxed{2334}_{5+4t}$

At $t=4$, all 6 reorderings are simultaneously normal ordered. In a sense three solitons collide all together at $t=4$. Compare this with the following time evolution pattern.

$t=0$: 11112222111133211431111111111111111111111111111111111
$t=1$: 11111111222211133214311111111111111111111111111111111
$t=2$: 11111111111222211332431111111111111111111111111111111
$t=3$: 11111111111111122221324331111111111111111111111111111
$t=4$: 11111111111111111112213224332111111111111111111111111
$t=5$: 11111111111111111111221132214332111111111111111111111
$t=6$: 11111111111111111111122111322114332111111111111111111
$t=7$: 11111111111111111111111221111322111433211111111111111
$t=8$: 11111111111111111111111112211111322111143321111111111
$t=9$: 11111111111111111111111111112211111132211114332111111

E.2. **Vertex operator construction of paths from scattering data.** Here we illustrate the action of the vertex operators $\Phi^{(1)},\dots,\Phi^{(n)}$ introduced in Section D.3 (D.4). It is convenient to use the vertex type diagram to express the action of the combinatorial R. For example the following successive actions of the combinatorial R

$$a\otimes b\otimes c\simeq b'\otimes a'\otimes c\simeq b'\otimes c'\otimes a'',$$

will be depicted by the diagram:



Given a path p and an element $b\in B_l$, one can carry b through p to the right by successively applying the combinatorial R as

$$b\otimes p\simeq p'\otimes b',\qquad p,p'\in B_{k_1}\otimes B_{k_2}\otimes\cdots\otimes B_{k_N}, \tag{E.2}$$

under the isomorphism $B_l\otimes(B_{k_1}\otimes\cdots\otimes B_{k_N})\simeq(B_{k_1}\otimes\cdots\otimes B_{k_N})\otimes B_l$. As the result we get $b'\in B_l$ and another path p'. Actually, the only situation $b'=u_l$ (highest element of B_l) will be encountered in our case, and the relation (E.2) will be denoted by $\Phi_b(p)=p'$. This is an elementary vertex operator. The previous ones $\Phi^{(1)},\dots,\Phi^{(n)}$ defined by (D.5) are compositions of Φ_b with several b.

For example, to calculate $\Phi_{\boxed{2334}}\left(\boxed{1}^{\otimes 5}\right)$, the relevant diagram is



Therefore we obtain $\Phi_{\boxed{2334}}\left(\boxed{1}^{\otimes 5}\right) = 43321$. Note that Φ_b has created one soliton labeled by the letters in b.

In general, if $b_1[d_1] \otimes \cdots \otimes b_m[d_m]$ is a normal ordered scattering data, $\Phi^{(1)}$ defined by (D.5) is realized as the following composition of elementary vertex operators:

$$\Phi^{(1)} = T_1^{d_1} \circ \Phi_{b_1} \circ T_1^{d_2-d_1} \circ \cdots \circ T_1^{d_m-d_{m-1}} \circ \Phi_{b_m}, \tag{E.3}$$

where $f \circ g(p) = f(g(p))$. The superscript "(1)" corresponds to that of $\mu^{(1)}$. Note that for $a = 1$, the effect of $\mathcal{T}_a^d$ in (D.5) is described by $T_1^d = (\Phi_{\boxed{1}})^d$.

In what follows we illustrate Theorem D.1 and Corollary D.2.

Example E.3. Take a path $p = 11112221322433$, which we have already considered in Example C.2 and Example E.1. From $t = 5$ case of Example E.1, the both sides of

$$\boxed{2222}_4 \otimes \boxed{23}_6 \otimes \boxed{334}_6 \simeq \boxed{2222}_4 \otimes \boxed{233}_6 \otimes \boxed{34}_6 \tag{E.4}$$

serve as the normal ordered scattering data $\mathcal{C}^{(1)}(p^{(1)})$. According to Theorem D.1 and (D.8), the original path p is reconstructed as $p = \Phi^{(1)}\mathcal{C}^{(1)}(p^{(1)})$. This $\Phi^{(1)}$ is realized, according to (E.3) and (E.4), as the following compositions of elementary vertex operators:

$$\begin{aligned} p &= T_1^4 \circ \Phi_{\boxed{2222}} \circ T_1^2 \circ \Phi_{\boxed{23}} \circ \Phi_{\boxed{334}} \left(\boxed{1}^{\otimes 14}\right) \\ &= T_1^4 \circ \Phi_{\boxed{2222}} \circ T_1^2 \circ \Phi_{\boxed{233}} \circ \Phi_{\boxed{34}} \left(\boxed{1}^{\otimes 14}\right). \end{aligned}$$

It is easy to check $p = 11112221322433$ from these formulas.

Let us illustrate Corollary D.2, which reflects the nested structure of the KKR bijection. For $\Phi^{(a)}$ (D.5) with general a, the formula (E.3) is replaced by

$$\Phi^{(a)} = (\Phi_{\boxed{a}})^{d_1} \circ \Phi_{b_1} \circ (\Phi_{\boxed{a}})^{d_2-d_1} \circ \cdots \circ (\Phi_{\boxed{a}})^{d_m-d_{m-1}} \circ \Phi_{b_m}. \tag{E.5}$$

Example E.4. We consider the same example as above. In the rigged configuration (see Example E.1), first look at the rightmost two diagrams which form an $A_1^{(1)}$ rigged configuration:



From $\mu^{(3)}$, we set $p^{(3)} = \boxed{4}$ according to (D.6) and obtain the scattering data $\mathcal{C}^{(3)}(p^{(3)}) = \boxed{4}_1$, which is obviously normal ordered. From (E.5), the A_1 highest path $p^{(2)} = \Phi^{(3)}\mathcal{C}^{(3)}(p^{(3)})$ with letters 3 and 4 is constructed as

$$p^{(2)} = \Phi_{\boxed{3}} \circ \Phi_{\boxed{4}} \left(\boxed{3} \otimes \boxed{333}\right) = \boxed{3} \otimes \boxed{334}.$$

Taking the rigging attached to $\mu^{(2)}$ into account, we obtain the normal ordered scattering data

$$\mathcal{C}^{(2)}(p^{(2)}) = \boxed{3}_1 \otimes \boxed{334}_4.$$

Next we look at the following parts



Then the A_2 highest path $p^{(1)} = \Phi^{(2)}\mathcal{C}^{(2)}(p^{(2)})$ with letters 2, 3 and 4 is calculated along (E.5) as

$$\begin{aligned} p^{(1)} &= \Phi_{\boxed{2}} \circ \Phi_{\boxed{3}} \circ (\Phi_{\boxed{2}})^3 \circ \Phi_{\boxed{334}} \left(\boxed{22} \otimes \boxed{222} \otimes \boxed{2222} \right) \\ &= \boxed{22} \otimes \boxed{223} \otimes \boxed{2334}. \end{aligned}$$

As a result, we have reproduced (E.1), which was the starting point of the previous Example E.3. Summarizing, the path $p = 11112221322433$ has been obtained as $p = \Phi^{(1)}C^{(1)}\Phi^{(2)}C^{(2)}\Phi^{(3)}C^{(3)}(p^{(3)})$.

References

[1] R. J. Baxter, *Exactly solved models in statistical mechanics*, Academic Press, London (1982).

[2] H. A. Bethe, Zur Theorie der Metalle, I. Eigenwerte und Eigenfunktionen der linearen Atomkette, Z. Physik **71** (1931) 205–231.

[3] M. Gaudin, *La fonction d'onde de Bethe*, Masson, Paris (1983).

[4] V. E. Korepin, N. M. Bogoliubov and A. G. Izergin, *Quantum Inverse Scattering Method and Correlation Functions*, Cambridge Univ. Press, (1997).

[5] M. Takahashi, *Thermodynamics of One-Dimensional Solvable Models*, Cambridge Univ. Press, (1999).

[6] G. E. Andrews, R. J. Baxter and P. J. Forrester, *Eight vertex SOS model and generalized Rogers-Ramanujan-type identities*, J. Stat. Phys. **35**, (1984) 193–266.

[7] E. Date, M. Jimbo, A. Kuniba, T. Miwa, M. Okado,: Exactly solvable SOS models: Local height probabilities and theta function identities, Nucl. Phys. B**290** [FS20] (1987) 231-273, II: Proof of the star-triangle relation and combinatorial identities, Adv. Stud. Pure Math. **16** (1988) 17-122.

[8] *Combinatorial Aspect of Integrable Systems*, eds. A. Kuniba and M. Okado, MSJ Memoirs **17** (2007).

[9] S. V. Kerov, A. N. Kirillov and N. Yu. Reshetikhin, Combinatorics, the Bethe ansatz and representations of the symmetric group. J. Soviet Math. **41** (1988) 916–924.

[10] A. N. Kirillov and N. Yu. Reshetikhin, The Bethe ansatz and the combinatorics of Young tableaux. J. Soviet Math. **41** (1988) 925–955.

[11] M. Kashiwara, On crystal bases of the q-analogue of universal enveloping algebras. Duke Math. J. **63** (1991) 465–516.

[12] S-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima and A. Nakayashiki, Affine crystals and vertex models, Int. J. Mod. Phys. A **7** (suppl. 1A), (1992) 449–484.

[13] A. Berkovich, B. M. McCoy and A. Schilling, Rogers-Schur-Ramanujan type identities for the $M(p, p')$ minimal models of conformal field theory, Commun.Math.Phys. **191** (1998) 325–395.

[14] S. Dasmahapatra, R. Kedem, T.R. Klassen, B.M. McCoy and E. Melzer, Quasi-Particles, Conformal Field Theory, and q-Series, Int. J. Mod. Phys. **B7** (1993) 3617–3648.

[15] B. L. Feigin and A. V. Stoyanovsky, Quasi-particle models for the representations of Lie algebras and geometry of flag manifold, (hep-th/9308079).

[16] O. Foda and T. A. Welsh, *Melzer's identities revisited*, Contemporary Math. **248** (1999) 207–234.

[17] A. Nakayashiki and Y. Yamada, Kostka polynomials and energy functions in solvable lattice models, Selecta Mathematica, New Ser. **3** (1997) 547–599.

[18] S. O. Warnaar, *Fermionic solution of the Andrews-Baxter-Forrester model I: unification of TBA and CTM methods*, J. Stat. Phys. **82** (1996) 657–685.

[19] G. Hatayama, A. Kuniba, M. Okado, T. Takagi and Y. Yamada, Remarks on fermionic formula, Contemp. Math. **248** (AMS 1999) 243–291.

[20] G. Hatayama, A. Kuniba, M. Okado, T. Takagi and Z. Tsuboi: Paths, Crystals and Fermionic Formulae, Prog. in Math. Phys. **23** (2002) 205–272.

[21] I. Macdonald, *Symmetric functions and Hall polynomials*, 2nd edition, Oxford Univ. Press, New York (1995).

[22] A. N. Kirillov, A. Schilling and M. Shimozono, A bijection between Littlewood-Richardson tableaux and rigged configurations. Selecta Math. **8** (2002) 67–135.

[23] A. Schilling, X=M Theorem: Fermionic formulas and rigged configurations under review, *Combinatorial Aspect in Integrable Systems*, MSJ Memoirs **17** (2007) 75–104.

[24] A. Schilling and M. Shimozono, X=M for symmetric powers, J. Alg. **295** (2006) 562–610.

[25] M. Okado, A. Schilling and M. Shimozono, A crystal to rigged configuration bijection for nonexceptional affine algebras, *Algebraic Combinatorics and Quantum Groups*, ed. N. Jing, World Scientific (2003), 85–124.

[26] G. Hatayama, K. Hikami, R. Inoue, A. Kuniba, T. Takagi and T. Tokihiro, The $A^{(1)}_M$ automata related to crystals of symmetric tensors, *J. Math. Phys.* **42** (2001) 274–308.

[27] K. Fukuda, M. Okado, Y. Yamada, Energy functions in box ball systems, Int. J. Mod. Phys. A **15** (2000) 1379–1392.
[28] D. Takahashi, On some soliton systems defined by using boxes and balls, Proceedings of the International Symposium on Nonlinear Theory and Its Applications (NOLTA '93), (1993) 555–558.
[29] D. Takahashi and J. Satsuma, A soliton cellular automaton, J. Phys. Soc. Jpn. **59** (1990) 3514–3519.
[30] G. Hatayama, A. Kuniba and T. Takagi: Soliton cellular automata associated with crystal bases, Nucl. Phys. B**577**[PM] (2000) 619–645.
[31] G. Hatayama, A. Kuniba, M. Okado, T. Takagi and Y. Yamada, Scattering rules in soliton cellular automata associated with crystal bases, *Contemporary Math.* **297** (2002) 151–182.
[32] A. Kuniba, M. Okado and Y. Yamada, Box-ball system with reflecting end, J. Nonlin. Math. Phys. **12** (2005) 475–507.
[33] T. Tokihiro, D. Takahashi, J. Matsukidaira and J. Satsuma, From soliton equations to integrable cellular automata through a limiting procedure, Phys. Rev. Lett. **76** (1996) 3247–3250.
[34] A. Kuniba, M. Okado, R. Sakamoto, T. Takagi and Y. Yamada: Crystal interpretation of Kerov-Kirillov-Reshetikhin bijection, Nucl. Phys. B**740** [PM] (2006) 299–327.
[35] R. Sakamoto, Crystal interpretation of Kerov-Kirillov-Reshetikhin bijection II. Proof for $\mathfrak{sl}_n$ Case, to appear in J. Algebraic Combinatorics, (math.QA/0601697).
[36] M. Sato and Y. Sato, Soliton Equations as Dynamical Systems on Infinite Dimensional Grassmann Manifold, *Nonlinear PDE in Applied Science*, U.S.-Japan Seminar, Tokyo, 1982, Lecture Notes in Num. Appl. Anal. **5** (1982) 259–271.
[37] M. Jimbo and T. Miwa, Solitons and infinite dimensional Lie algebras, Publ. RIMS. Kyoto Univ. **19** (1983) 943–1001.
[38] Y. Yamada, A birational representation of Weyl group, combinatorial R-matrix and discrete Toda equation, in "Physics and Combinatorics 2000", eds. A. N. Kirillov and N. Liskova (World Scientific, 2001) 305–319.
[39] J. S. Birman: *Braids, Links, and Mapping Class Groups*, Princeton University Press (1974).
[40] L. Deka and A. Schilling, New fermionic formula for unrestricted Kostka polynomials, J. Combinatorial Theory, Series A **113** (2006) 1435–1461.
[41] C. L. Schultz, Eigenvectors of the multicomponent generalization of the six-vertex model, Physica. **A122** (1983) 71–88.

Atsuo Kuniba:
Institute of Physics, Graduate School of Arts and Sciences, University of Tokyo, Komaba, Tokyo 153-8902, Japan
atsuo@gokutan.c.u-tokyo.ac.jp

Reiho Sakamoto:
Department of Physics, Graduate School of Science, University of Tokyo, Hongo, Tokyo 113-0033, Japan
reiho@monet.phys.s.u-tokyo.ac.jp

Yasuhiko Yamada:
Department of Mathematics, Faculty of Science, Kobe University, Hyogo 657-8501, Japan
yamaday@math.kobe-u.ac.jp